

ZNF16 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17131a

Specification

ZNF16 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P17020
Other Accession	NP_001025147.2 , NP_008889.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	76472
Antigen Region	53-81

ZNF16 Antibody (N-term) - Additional Information

Gene ID 7564

Other Names

Zinc finger protein 16, Zinc finger protein KOX9, ZNF16, HZF1, KOX9

Target/Specificity

This ZNF16 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 53-81 amino acids from the N-terminal region of human ZNF16.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ZNF16 Antibody (N-term) is for research use only and not for use in diagnostic or



ZNF16 Antibody (N-term) (Cat. #AP17131a) western blot analysis in MDA-MB453 cell line lysates (35ug/lane). This demonstrates the ZNF16 antibody detected the ZNF16 protein (arrow).

ZNF16 Antibody (N-term) - Background

The protein encoded by this gene contains a C2H2 type of zinc finger, and thus may function as a transcription factor. This gene is located in a region close to ZNF7/KOX4, a gene also encoding a zinc finger protein, on chromosome 8. Two alternatively spliced variants, encoding the same protein, have been identified.

ZNF16 Antibody (N-term) - References

- Dreier, B., et al. J. Mol. Biol. 303(4):489-502(2000)
- Rousseau-Merck, M.F., et al. Ann. Genet. 38(2):81-84(1995)
- Thiesen, H.J. New Biol. 2(4):363-374(1990)

therapeutic procedures.

ZNF16 Antibody (N-term) - Protein Information

Name ZNF16

Synonyms HZF1, KOX9

Function

Acts as a transcriptional activator. Promotes cell proliferation by facilitating the cell cycle phase transition from the S to G2/M phase. Involved in both the hemin- and phorbol myristate acetate (PMA)-induced erythroid and megakaryocytic differentiation, respectively. Plays also a role as an inhibitor of cell apoptosis.

Cellular Location

Nucleus.

Tissue Location

Ubiquitous..

ZNF16 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)