

Bmp3 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1714a

Specification

Bmp3 Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P12645
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	53372
Antigen Region	8-36

Bmp3 Antibody (N-term) - Additional Information

Gene ID 651

Other Names

Bone morphogenetic protein 3, BMP-3, Bone morphogenetic protein 3A, BMP-3A, Osteogenin, BMP3, BMP3A

Target/Specificity

This Bmp3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 8-36 amino acids from the N-terminal region of human Bmp3.

Dilution

WB~~1:1000
 IHC-P~~1:50~100

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Bmp3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.



Western blot analysis of hBmp3-E23 (Cat. #AP1714a) in 293 cell line lysates (35ug/lane). BMP3 (arrow) was detected using the purified Pab.



Western blot analysis of Bmp3 (arrow) using rabbit polyclonal Bmp3 Antibody (N-term) (Cat.#AP1714a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the Bmp3 gene (Lane 2) (Origene Technologies).

Bmp3 Antibody (N-term) - Protein Information**Name** BMP3**Synonyms** BMP3A**Function**

Growth factor of the TGF-beta superfamily that plays an essential role in developmental process by inducing and patterning early skeletal formation and by negatively regulating bone density.

Antagonizes the ability of certain osteogenic BMPs to induce osteoprogenitor differentiation and ossification (PubMed:11138004, PubMed:15269709). Initiates signaling cascades by associating with type II receptor ACVR2B to activate SMAD2-dependent and SMAD-independent signaling cascades including TAK1 and JNK pathways (PubMed:31665064).

Cellular Location

Secreted.


Tissue Location

Expressed in adult and fetal cartilage.

Bmp3 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Bmp3 Antibody (N-term) - Background

The BMPs belong to the TGF-Beta superfamily. BMPs stimulate the production of bone matrix proteins and thereby alter stromal cell and osteoclast proliferation, a key step in bone marrow development. In addition to promoting ectopic bone and cartilage development, BMPs regulate the growth, differentiation, chemotaxis, proliferation, and apoptosis of various cell types, including mesenchymal cells, epithelial cells, hematopoietic cells, and neuronal cells. BMPs also play a role in dorsal/ventral patterning. BMP3, highly expressed in lung, ovary and small intestine, participates in cartilage and bone formation. BMP3 and BMP2 genes map to conserved regions between human and mouse.

Bmp3 Antibody (N-term) - References

Jin, Y., et al., *Histol. Histopathol.* 16(4):1013-1019 (2001).
Tabas, J.A., et al., *Genomics* 9(2):283-289 (1991).
Dickinson, M.E., et al., *Genomics* 6(3):505-520 (1990).
Wozney, J.M., et al., *Science* 242(4885):1528-1534 (1988).