

Bmp6 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1717A

Specification

Bmp6 Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P22004
Other Accession	NP_001709
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	25-55

Bmp6 Antibody (N-term) - Additional Information

Gene ID 654

Other Names

Bone morphogenetic protein 6, BMP-6, VG-1-related protein, VG-1-R, VGR-1, BMP6, VGR

Target/Specificity

This Bmp6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 25-55 amino acids from the N-terminal region of human Bmp6.

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Bmp6 Antibody (N-term) is for research use only and not for use in diagnostic or



The anti-Bmp6 (N-term) Pab (Cat. #AP1717a) is used in Western blot to detect Bmp6 in HL60 cell lysate.



Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with Bmp6 antibody (N-term)(Cat. #AP1717a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

therapeutic procedures.

Bmp6 Antibody (N-term) - Protein Information

Name BMP6

Synonyms VGR

Function

Growth factor of the TGF-beta superfamily that plays essential roles in many developmental processes including cartilage and bone formation (PubMed:31019025). Plays also an important role in the regulation of iron metabolism by acting as a ligand for hemojuvelin/HJV (By similarity). Initiates the canonical BMP signaling cascade by associating with type I receptor ACVR1 and type II receptor ACVR2B (PubMed:18070108). In turn, ACVR1 propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressors of transcription of target. Can also signal through non- canonical pathway such as TAZ-Hippo signaling cascade to modulate VEGF signaling by regulating VEGFR2 expression (PubMed:33021694).

Cellular Location

Secreted.

Bmp6 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Bmp6 Antibody (N-term) - Citations

- [Increased BMP6 levels in the brains of Alzheimer's disease patients and APP transgenic](#)

Bmp6 Antibody (N-term) - Background

The bone morphogenetic proteins (BMPs) are a family of secreted signaling molecules that can induce ectopic bone growth. Many BMPs are part of the transforming growth factor-beta (TGFB) superfamily. BMPs were originally identified by an ability of demineralized bone extract to induce endochondral osteogenesis in vivo in an extraskeletal site. Based on its expression early in embryogenesis, Bmp6 has a proposed role in early development. In addition, the fact that BMP6 is closely related to BMP5 and BMP7 has lead to speculation of possible bone inductive activity.

Bmp6 Antibody (N-term) - References

Lories, R.J., et al., *Arthritis Rheum.* 48(10):2807-2818 (2003).
Bobacz, K., et al., *Arthritis Rheum.* 48(9):2501-2508 (2003).
Tamada, H., et al., *Biochim. Biophys. Acta* 1395(3):247-251 (1998).
Rickard, D.J., et al., *J. Clin. Invest.* 101(2):413-422 (1998).
Olavesen, M.G., et al., *Genomics* 46(2):303-306 (1997).

mice are accompanied by impaired neurogenesis.