

PJA2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17220c

Specification

PJA2 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O43164
Other Accession	NP_055634.3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	78214
Antigen Region	284-311

PJA2 Antibody (Center) - Additional Information

Gene ID 9867

Other Names

E3 ubiquitin-protein ligase Praja-2, Praja2, 632-, RING finger protein 131, PJA2, KIAA0438, RNF131

Target/Specificity

This PJA2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 284-311 amino acids from the Central region of human PJA2.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PJA2 Antibody (Center) is for research use only and not for use in diagnostic or



PJA2 Antibody (Center) (Cat. #AP17220c) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the PJA2 antibody detected the PJA2 protein (arrow).

PJA2 Antibody (Center) - Background

PJA2 has E2-dependent E3 ubiquitin-protein ligase activity (By similarity).

PJA2 Antibody (Center) - References

Olsen, J.V., et al. Cell 127(3):635-648(2006)
Sasaki, A., et al. J. Biol. Chem. 277(25):22541-22546(2002)
Yu, P., et al. Genomics 79(6):869-874(2002)
Mishra, L., et al. Oncogene 15(19):2361-2368(1997)
Nakayama, M., et al. J. Neurosci. 15 (7 PT 2), 5238-5248 (1995) :

therapeutic procedures.

PJA2 Antibody (Center) - Protein Information

Name PJA2

Synonyms KIAA0438, RNF131

Function

Has E2-dependent E3 ubiquitin-protein ligase activity. Responsible for ubiquitination of cAMP-dependent protein kinase type I and type II-alpha/beta regulatory subunits and for targeting them for proteasomal degradation. Essential for PKA-mediated long-term memory processes. Through the ubiquitination of MFHAS1, positively regulates the TLR2 signaling pathway that leads to the activation of the downstream p38 and JNK MAP kinases and promotes the polarization of macrophages toward the pro-inflammatory M1 phenotype (PubMed:28471450).

Cellular Location

Cytoplasm. Cell membrane. Endoplasmic reticulum membrane; Peripheral membrane protein. Golgi apparatus membrane; Peripheral membrane protein. Cell junction, synapse Cell junction, synapse, postsynaptic density Note=Localizes at the cytoplasmic side of endoplasmic reticulum and Golgi apparatus. Expressed in the postsynaptic density region of synapses (By similarity). Colocalizes with PRKAR2A and PRKAR2B in the cytoplasm and the cell membrane.

PJA2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)