

IER5 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17351C

Specification

IER5 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q5VY09
Other Accession	NP_057629.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	33704
Antigen Region	122-149

IER5 Antibody (Center) - Additional Information

Gene ID 51278

Other Names

Immediate early response gene 5 protein,
IER5

Target/Specificity

This IER5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 122-149 amino acids from the Central region of human IER5.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IER5 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.



IER5 Antibody (Center) (Cat. #AP17351c) western blot analysis in MCF-7 cell line lysates (35ug/lane). This demonstrates the IER5 antibody detected the IER5 protein (arrow).

IER5 Antibody (Center) - Background

This gene encodes a protein that is similar to other immediate early response proteins. In the mouse, a similar gene may play an important role in mediating the cellular response to mitogenic signals. Studies in rats found the expression of a similar gene to be increased after waking and sleep deprivation.

IER5 Antibody (Center) - References

Ding, K.K., et al. Radiat Environ Biophys 48(2):205-213(2009)
Wan, D., et al. Proc. Natl. Acad. Sci. U.S.A. 101(44):15724-15729(2004)
Cirelli, C., et al. Brain Res. 885(2):303-321(2000)
Williams, M., et al. Genomics 55(3):327-334(1999)

IER5 Antibody (Center) - Protein Information**Name** IER5**Function**

Plays a role as a transcription factor (PubMed:22132193, PubMed:25355627). Mediates positive transcriptional regulation of several chaperone genes during the heat shock response in a HSF1- dependent manner (PubMed:25355627, PubMed:25816751). Mediates negative transcriptional regulation of CDC25B expression (PubMed:22132193). Plays a role in the dephosphorylation of the heat shock factor HSF1 and ribosomal protein S6 kinase (S6K) by the protein phosphatase PP2A (PubMed:25816751, PubMed:26496226). Involved in the regulation of cell proliferation and resistance to thermal stress (PubMed:22132193, PubMed:25355627, PubMed:26496226). Involved in the cell cycle checkpoint and survival in response to ionizing radiation (PubMed:19238419, PubMed:22132193). Associates with chromatin to the CDC25B promoter (PubMed:22132193).

Cellular Location

Nucleus. Cytoplasm. Note=Predominantly cytoplasmic (PubMed:25816751).

Translocated in the nucleus during heat shock (PubMed:25816751).

Tissue Location

Expressed in acute myeloid leukemia (AML) cells.

IER5 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IER5 Antibody (Center) - Citations

- [IER family proteins are regulators of protein phosphatase PP2A and modulate the phosphorylation status of CDC25A.](#)
- [Regulation of the stability and activity of CDC25A and CDC25B by protein phosphatase PP2A and 14-3-3 binding.](#)