

TET2 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17372a

Specification

TET2 Antibody (N-term) - Product Information

| | |
|-------------------|---|
| Application | WB, IHC-P,E |
| Primary Accession | Q6N021 |
| Other Accession | NP_001120680.1 , NP_060098.3 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit Ig |
| Calculated MW | 223811 |
| Antigen Region | 528-554 |

TET2 Antibody (N-term) - Additional Information

Gene ID 54790

Other Names

Methylcytosine dioxygenase TET2,
11411n2, TET2, KIAA1546

Target/Specificity

This TET2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 528-554 amino acids from the N-terminal region of human TET2.

Dilution

WB~~1:2000
IHC-P~~1:100

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TET2 Antibody (N-term) is for research use



Anti-TET2 Antibody (N-term) at 1:2000 dilution + MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 224 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of AP17372a on paraffin-embedded Human placenta tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope

only and not for use in diagnostic or therapeutic procedures.

TET2 Antibody (N-term) - Protein Information

Name TET2

Synonyms KIAA1546

Function

Dioxygenase that catalyzes the conversion of the modified genomic base 5-methylcytosine (5mC) into 5-hydroxymethylcytosine (5hmC) and plays a key role in active DNA demethylation. Has a preference for 5-hydroxymethylcytosine in CpG motifs. Also mediates subsequent conversion of 5hmC into 5-formylcytosine (5fC), and conversion of 5fC to 5-carboxylcytosine (5caC). Conversion of 5mC into 5hmC, 5fC and 5caC probably constitutes the first step in cytosine demethylation. Methylation at the C5 position of cytosine bases is an epigenetic modification of the mammalian genome which plays an important role in transcriptional regulation. In addition to its role in DNA demethylation, also involved in the recruitment of the O-GlcNAc transferase OGT to CpG-rich transcription start sites of active genes, thereby promoting histone H2B GlcNAcylation by OGT.

Tissue Location

Broadly expressed. Highly expressed in hematopoietic cells; highest expression observed in granulocytes. Expression is reduced in granulocytes from peripheral blood of patients affected by myelodysplastic syndromes.

retrieval was performed by EDTA buffer (pH9.0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

TET2 Antibody (N-term) - Background

TET2 may catalyze the conversion of methylcytosine (5mC) to 5-hydroxymethylcytosine (hmC) (By similarity). Play an important role in the regulation of myelopoiesis.

TET2 Antibody (N-term) - References

Mallo, M., et al. Haematologica 95(10):1798-1800(2010)
Smith, A.E., et al. Blood (2010) In press :
Tefferi, A., et al. Leukemia 24(7):1302-1309(2010)
Kim, S.T., et al. Prostate (2010) In press :
Rocquain, J., et al. BMC Cancer 10, 401 (2010) :

TET2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)