

NOSIP Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17578a

Specification

NOSIP Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q9Y314
Other Accession	Q4R7H4 , NP_057037.1
Reactivity	Human
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	33172
Antigen Region	67-95

NOSIP Antibody (N-term) - Additional Information

Gene ID 51070

Other Names

Nitric oxide synthase-interacting protein,
eNOS-interacting protein, NOSIP

Target/Specificity

This NOSIP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 67-95 amino acids from the N-terminal region of human NOSIP.

Dilution

WB~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NOSIP Antibody (N-term) is for research use



NOSIP Antibody (N-term) (Cat. #AP17578a) western blot analysis in Ramos cell line lysates (35ug/lane). This demonstrates the NOSIP antibody detected the NOSIP protein (arrow).

NOSIP Antibody (N-term) - Background

NOSIP negatively regulates nitric oxide production by inducing NOS1 and NOS3 translocation to actin cytoskeleton and inhibiting their enzymatic activity.

NOSIP Antibody (N-term) - References

- Matsuoka, S., et al. Science 316(5828):1160-1166(2007)
- Lamesch, P., et al. Genomics 89(3):307-315(2007)
- Schleicher, M., et al. Mol. Cell. Biol. 25(18):8251-8258(2005)
- Andersen, J.S., et al. Nature 433(7021):77-83(2005)
- Dreyer, J., et al. J. Neurosci. 24(46):10454-10465(2004)

only and not for use in diagnostic or therapeutic procedures.

NOSIP Antibody (N-term) - Protein Information

Name NOSIP

Function

E3 ubiquitin-protein ligase that is essential for proper development of the forebrain, the eye, and the face. Catalyzes monoubiquitination of serine/threonine-protein phosphatase 2A (PP2A) catalytic subunit PPP2CA/PPP2CB (By similarity). Negatively regulates nitric oxide production by inducing NOS1 and NOS3 translocation to actin cytoskeleton and inhibiting their enzymatic activity (PubMed:<<http://www.uniprot.org/citations/11149895>>, PubMed:<<http://www.uniprot.org/citations/15548660>>, PubMed:<<http://www.uniprot.org/citations/16135813>>).

Cellular Location

Cytoplasm. Nucleus. Note=Translocates from nucleus to cytoplasm in the G2 phase of the cell cycle (PubMed:16135813)

Tissue Location

Expressed in heart, brain and lung. Present in endothelial cells (at protein level).

NOSIP Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)