

### **FANCB Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP18051C

## **Specification**

#### **FANCB Antibody (Center) - Product Information**

Application WB,E
Primary Accession Q8NB91

Other Accession NP 001018123.1

Reactivity
Host
Clonality
Isotype
Calculated MW
Antigen Region

Human
Rabbit
Polyclonal
Rabbit Ig
97726
360-388

**FANCB Antibody (Center) - Additional Information** 

#### **Gene ID 2187**

#### **Other Names**

Fanconi anemia group B protein, Protein FACB, Fanconi anemia-associated polypeptide of 95 kDa, FAAP95, FANCB

## Target/Specificity

This FANCB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 360-388 amino acids from the Central region of human FANCB.

## **Dilution**

WB~~1:1000

### **Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

## Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

FANCB Antibody (Center) is for research use only and not for use in diagnostic or



FANCB Antibody (Center) (Cat. #AP18051c) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the FANCB antibody detected the FANCB protein (arrow).

## **FANCB Antibody (Center) - Background**

The Fanconi anemia complementation group (FANC) currently

includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2,

FANCE, FANCF, FANCG, FANCI, FANCJ (also called BRIP1), FANCL, FANCM

and FANCN (also called PALB2). The previously defined group FANCH

is the same as FANCA. Fanconi anemia is a genetically heterogeneous

recessive disorder characterized by

cytogenetic instability, hypersensitivity to DNA crosslinking agents,

increased chromosomal


breakage, and defective DNA repair. The members of the Fanconi

anemia complementation group do not share sequence similarity; they

are related by their assembly into a common nuclear protein

complex. This gene encodes the protein for complementation group B.

Alternative splicing results in two transcript variants encoding





Tel: 858.875.1900 Fax: 858.622.0609

therapeutic procedures.

FANCB Antibody (Center) - Protein Information

**Name FANCB** 

**Function** 

DNA repair protein required for FANCD2 ubiquitination.

**Cellular Location** Nucleus.

# **FANCB Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

the same protein.

12(10):1111-1122(2007)

# **FANCB Antibody (Center) - References**

Cheung, K.L., et al. World J Surg Oncol 8, 38 (2010): Smith, I.M., et al. ORL J. Otorhinolaryngol. Relat. Spec. 72(1):44-50(2010) Barroso, E., et al. Breast Cancer Res. Treat. 118(3):655-660(2009) Garcia, M.J., et al. Breast Cancer Res. Treat. 113(3):545-551(2009) Nomura, Y., et al. Genes Cells