

LAMP2 Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1824d

Specification

LAMP2 Antibody - Product Information

Application	WB,E
Primary Accession	P13473
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig

LAMP2 Antibody - Additional Information

Gene ID 3920

Other Names

Lysosome-associated membrane glycoprotein 2, LAMP-2, Lysosome-associated membrane protein 2, CD107 antigen-like family member B, CD107b, LAMP2

Target/Specificity

This LAMP2 antibody is generated from rabbits immunized with LAMP2 recombinant protein.

Dilution

WB~1:2000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

LAMP2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



All lanes : Anti-LAMP2 Antibody at 1:1000 dilution Lane 1: Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Observed band size : 120kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-LAMP2 Antibody at 1:2000 dilution Lane 1:HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),


LAMP2 Antibody - Protein Information

Name LAMP2

Function

Plays an important role in chaperone-mediated autophagy, a process that mediates lysosomal degradation of proteins in response to various stresses and as part of the normal turnover of proteins with a long biological half-life (PubMed:8662539, PubMed:11082038, PubMed:18644871, PubMed:24880125, PubMed:27628032). Functions by binding target proteins, such as GAPDH and MLLT11, and targeting them for lysosomal degradation (PubMed:8662539, PubMed:11082038, PubMed:18644871, PubMed:24880125). Plays a role in lysosomal protein degradation in response to starvation (By similarity). Required for the fusion of autophagosomes with lysosomes during autophagy (PubMed:27628032). Cells that lack LAMP2 express normal levels of VAMP8, but fail to accumulate STX17 on autophagosomes, which is the most likely explanation for the lack of fusion between autophagosomes and lysosomes (PubMed:27628032). Required for normal degradation of the contents of autophagosomes (PubMed:27628032).

Peroxidase conjugated at 1/10000 dilution.
Observed band size : 110kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-LAMP2 Antibody at 1:2000 dilution
Lane 1: H. placenta whole lysate
Lane 2: HeLa whole cell lysate
Lane 3: HepG2 whole cell lysate
Lane 4: SK-MEL-2 whole cell lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 45 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

LAMP2 Antibody - Background

LAMP2 is a member of a family of membrane glycoproteins. This glycoprotein provides selectins with carbohydrate ligands. It may play a role in tumor cell metastasis. It may also function in the protection, maintenance, and adhesion of the lysosome.

LAMP2 Antibody - References

- Sarafian,V.S., Acta. Biol. Hung. 57 (3), 315-322 (2006)
- Liu,T., J. Proteome Res. 4 (6), 2070-2080 (2005)
- Mane,S.M., Arch. Biochem. Biophys. 268 (1), 360-378 (1989)
- Fukuda,M., J. Biol. Chem. 263 (35), 18920-18928 (1988)

target="_blank">27628032). Required for efficient MHCII-mediated presentation of exogenous antigens via its function in lysosomal protein degradation; antigenic peptides generated by proteases in the endosomal/lysosomal compartment are captured by nascent MHCII subunits (PubMed:20518820). Is not required for efficient MHCII-mediated presentation of endogenous antigens (PubMed:20518820).

Cellular Location

Cell membrane; Single-pass type I membrane protein
{ECO:0000255|PROSITE-ProRule:PRU00740, ECO:0000269|PubMed:17897319}
Endosome membrane; Single-pass type I membrane protein
{ECO:0000255|PROSITE-ProRule:PRU00740, ECO:0000269|PubMed:17897319}.
Lysosome membrane
{ECO:0000255|PROSITE-ProRule:PRU00740, ECO:0000269|PubMed:11082038, ECO:0000269|PubMed:17897319, ECO:0000269|PubMed:18644871, ECO:0000269|PubMed:2912382};
Single-pass type I membrane protein
{ECO:0000255|PROSITE-ProRule:PRU00740, ECO:0000269|PubMed:17897319}
Cytoplasmic vesicle, autophagosome membrane
{ECO:0000250|UniProtKB:P17047}.
Note=This protein shuttles between lysosomes, endosomes, and the plasma membrane

Tissue Location

Isoform LAMP-2A is highly expressed in placenta, lung and liver, less in kidney and pancreas, low in brain and skeletal muscle (PubMed:7488019, PubMed:26856698).
Isoform LAMP-2B is detected in spleen, thymus, prostate, testis, small intestine, colon, skeletal muscle, brain, placenta, lung, kidney, ovary and pancreas and liver (PubMed:7488019, PubMed:26856698).
Isoform LAMP-2C is detected in small intestine, colon, heart, brain, skeletal muscle, and at lower levels in kidney and placenta (PubMed:26856698).

LAMP2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

LAMP2 Antibody - Citations

- [Protein-carbohydrate ingestion alters Vps34 cellular localization independent of changes in kinase activity in human skeletal muscle](#)
- [Differential localisation and anabolic responsiveness of mTOR complexes in human skeletal muscle in response to feeding and exercise.](#)