

PPP4R2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18485c

Specification

PPP4R2 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O9NY27
Other Accession	NP_777567.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	46898
Antigen Region	228-254

PPP4R2 Antibody (Center) - Additional Information

Gene ID 151987

Other Names

Serine/threonine-protein phosphatase 4 regulatory subunit 2, PPP4R2

Target/Specificity

This PPP4R2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 228-254 amino acids from the Central region of human PPP4R2.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PPP4R2 Antibody (Center) is for research use only and not for use in diagnostic or



PPP4R2 Antibody (Center) - Background

Regulatory subunit of serine/threonine-protein phosphatase 4 (PP4). May regulate the activity of PPP4C at centrosomal microtubule organizing centers. Its interaction with the SMN complex leads to enhance the temporal localization of snRNPs, suggesting a role of PPP4C in maturation of spliceosomal snRNPs. The PPP4C-PPP4R2-PPP4R3A PP4 complex specifically dephosphorylates H2AFX phosphorylated on 'Ser-140' (gamma-H2AFX) generated during DNA replication and required for DNA DSB repair.

PPP4R2 Antibody (Center) - References

- Lee, D.H., et al. Nat. Struct. Mol. Biol. 17(3):365-372(2010)
- Chen, G.I., et al. J. Biol. Chem. 283(43):29273-29284(2008)
- Chowdhury, D., et al. Mol. Cell 31(1):33-46(2008)
- Ewing, R.M., et al. Mol. Syst. Biol. 3, 89 (2007)
- Gingras, A.C., et al. Mol. Cell Proteomics

therapeutic procedures.

4(11):1725-1740(2005)

PPP4R2 Antibody (Center) - Protein Information

Name PPP4R2

Function

Regulatory subunit of serine/threonine-protein phosphatase 4 (PP4). May regulate the activity of PPP4C at centrosomal microtubule organizing centers. Its interaction with the SMN complex leads to enhance the temporal localization of snRNPs, suggesting a role of PPP4C in maturation of spliceosomal snRNPs. The PPP4C-PPP4R2-PPP4R3A PP4 complex specifically dephosphorylates H2AX phosphorylated on 'Ser-140' (gamma-H2AX) generated during DNA replication and required for DNA double strand break repair. Mediates RPA2 dephosphorylation by recruiting PPP4C to RPA2 in a DNA damage-dependent manner. RPA2 dephosphorylation is required for the efficient RPA2-mediated recruitment of RAD51 to chromatin following double strand breaks, an essential step for DNA repair.

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Note=Ionizing radiation induces relocalization to nuclear foci and colocalization with RPA2

Tissue Location

Widely expressed..

PPP4R2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)