

NAAA Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18542b

Specification

NAAA Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q02083
Other Accession	NP_001035861.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	40066
Antigen Region	331-359

NAAA Antibody (C-term) - Additional Information

Gene ID 27163

Other Names

N-acylethanolamine-hydrolyzing acid amidase, 351-, Acid ceramidase-like protein, N-acylsphingosine amidohydrolase-like, ASAH-like protein, N-acylethanolamine-hydrolyzing acid amidase subunit alpha, N-acylethanolamine-hydrolyzing acid amidase subunit beta, NAAA, AS AHL, PLT

Target/Specificity

This NAAA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 331-359 amino acids from the C-terminal region of human NAAA.

Dilution


WB~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw



NAAA Antibody (C-term) (Cat. #AP18542b) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the NAAA antibody detected the NAAA protein (arrow).

NAAA Antibody (C-term) - Background

This gene encodes an N-acylethanolamine-hydrolyzing enzyme which is highly similar to acid ceramidase. Multiple transcript variants encoding different isoforms have been found for this gene.

NAAA Antibody (C-term) - References

- Wang, J., et al. J. Biochem. 144(5):685-690(2008)
- Muller, T.D., et al. Child Adolesc Psychiatry Ment Health 2 (1), 33 (2008) :
- Schulze, H., et al. Biol. Chem. 388(12):1333-1343(2007)
- Zhao, L.Y., et al. Biochim. Biophys. Acta 1771(11):1397-1405(2007)
- Oh, J.H., et al. Mamm. Genome 16(12):942-954(2005)

cycles.

Precautions

NAAA Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NAAA Antibody (C-term) - Protein Information

Name NAAA

{ECO:0000303|PubMed:30301806,
ECO:0000312|HGNC:HGNC:736}

Function

Degrades bioactive fatty acid amides to their corresponding acids, with the following preference: N-palmitoylethanolamine > N-myristoylethanolamine > N-lauroylethanolamine = N-stearoylethanolamine > N-arachidonylethanolamine > N-oleoylethanolamine (PubMed:15655246, PubMed:17980170, PubMed:18793752, PubMed:30301806, PubMed:22825852). Also exhibits weak hydrolytic activity against the ceramides N-lauroylsphingosine and N-palmitoylsphingosine (PubMed:15655246).

Cellular Location

Lysosome. Membrane; Peripheral membrane protein

Tissue Location

Expressed in numerous tissues, with highest levels in liver and kidney, followed by pancreas

NAAA Antibody (C-term) - Protocols

Provided below are standard protocols that you

may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)