

GNAS Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18552b

Specification

GNAS Antibody (C-term) - Product Information

Application	WB, E
Primary Accession	O95467
Other Accession	NP_057676.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	28029
Antigen Region	217-244

GNAS Antibody (C-term) - Additional Information

Gene ID 2778

Other Names

Neuroendocrine secretory protein 55, NESP55, LHAL tetrapeptide, GPIPIRRH peptide, GNAS (<u href="http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=4392" target="_blank">HGNC:4392</u>)

Target/Specificity

This GNAS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 217-244 amino acids from the C-terminal region of human GNAS.

Dilution


WB~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



GNAS Antibody (C-term) (Cat. #AP18552b) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the GNAS antibody detected the GNAS protein (arrow).

GNAS Antibody (C-term) - Background

This locus has a highly complex imprinted expression pattern. It gives rise to maternally, paternally, and biallelically expressed transcripts that are derived from four alternative promoters and 5' exons. Some transcripts contain a differentially methylated region (DMR) at their 5' exons, and this DMR is commonly found in imprinted genes and correlates with transcript expression. An antisense transcript is produced from an overlapping locus on the opposite strand. One of the transcripts produced from this locus, and the antisense transcript, are paternally expressed noncoding RNAs, and may regulate imprinting in this region. In addition, one of the transcripts contains a second overlapping ORF, which encodes a structurally unrelated protein

Precautions

GNAS Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

GNAS Antibody (C-term) - Protein Information

Name GNAS ([HGNC:4392](#))

Cellular Location

Cytoplasmic vesicle, secretory vesicle. Secreted. Note=Neuroendocrine secretory granules.

GNAS Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

- Alex. Alternative splicing of downstream exons is also observed, which results in different forms of the stimulatory G-protein alpha subunit, a key element of the classical signal transduction pathway linking receptor-ligand interactions with the activation of adenylyl cyclase and a variety of cellular responses. Multiple transcript variants encoding different isoforms have been found for this gene. Mutations in this gene result in pseudohypoparathyroidism type 1a, pseudohypoparathyroidism type 1b, Albright hereditary osteodystrophy, pseudopseudohypoparathyroidism, McCune-Albright syndrome, progressive osseus heteroplasia, polyostotic fibrous dysplasia of bone, and some pituitary tumors.

GNAS Antibody (C-term) - References

Idziaszczyk, S., et al. Cancer Genet. Cytogenet. 202(1):67-69(2010)
Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Tominaga, E., et al. Gynecol. Oncol. 118(2):160-166(2010)
Park, C.H., et al. Ann. Clin. Lab. Sci. 40(3):261-266(2010)
Cross, D.S., et al. BMC Genet. 11, 51 (2010) :