

JUN Antibody (Center T93)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18924C

Specification

JUN Antibody (Center T93) - Product Information

Application	IF, WB,E
Primary Accession	P05412
Other Accession	P17325 , P56432 , P05627 , O77627 , NP_002219.1
Reactivity Predicted	Human Bovine, Mouse, Pig, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	35676
Antigen Region	72-98

JUN Antibody (Center T93) - Additional Information

Gene ID 3725

Other Names

Transcription factor AP-1, Activator protein 1, AP1, Proto-oncogene c-Jun, V-jun avian sarcoma virus 17 oncogene homolog, p39, JUN

Target/Specificity

This JUN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 72-98 amino acids from the Central region of human JUN.

Dilution


IF~~1:10~50
WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C



Fluorescent confocal image of U251 cell stained with JUN Antibody (Center T93)(Cat#AP18924c).U251 cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with JUN primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C).Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7units/ml, 1 h at 37°C). Nuclei were counterstained with DAPI (blue) (10 µg/ml, 10 min). JUN immunoreactivity is localized to Nucleus significantly.



in small aliquots to prevent freeze-thaw cycles.

Precautions

JUN Antibody (Center T93) is for research use only and not for use in diagnostic or therapeutic procedures.

JUN Antibody (Center T93) - Protein Information

Name JUN

Function

Transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3' (PubMed:10995748, PubMed:22083952). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:17210646). Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:24623306). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:24623306).

Cellular Location

Nucleus.

Tissue Location

Expressed in the developing and adult prostate and prostate cancer cells.

JUN Antibody (Center T93) (Cat. #AP18924c) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the JUN antibody detected the JUN protein (arrow).

JUN Antibody (Center T93) - Background

This gene is the putative transforming gene of avian sarcoma virus 17. It encodes a protein which is highly similar to the viral protein, and which interacts directly with specific target DNA sequences to regulate gene expression. This gene is intronless and is mapped to 1p32-p31, a chromosomal region involved in both translocations and deletions in human malignancies.

JUN Antibody (Center T93) - References

Gonsalves, C., et al. J. Immunol. 185(10):6253-6264(2010)
Bozec, A., et al. J. Cell Biol. 190(6):1093-1106(2010)
Machida, K., et al. Hepatology 52(2):480-492(2010)
Madi, A., et al. BMC Microbiol. 10, 215 (2010) :
Johnatty, S.E., et al. PLoS Genet. 6 (7), E1001016 (2010) :

JUN Antibody (Center T93) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)