

Mouse Mcm2 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19218B

Specification

Mouse Mcm2 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P97310
Other Accession	NP_032590.2
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	102078
Antigen Region	713-742

Mouse Mcm2 Antibody (C-term) - Additional Information

Gene ID 17216

Other Names

DNA replication licensing factor MCM2,
 Minichromosome maintenance protein 2
 homolog, Nuclear protein BM28, Mcm2,
 Bm28, Cdcl1, Kiaa0030, Mcmd2

Target/Specificity

This Mouse Mcm2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 713-742 amino acids from the C-terminal region of mouse Mcm2.

Dilution


WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



Mouse Mcm2 Antibody (C-term) (Cat. #AP19218b) western blot analysis in mouse Neuro-2a cell line lysates (35ug/lane). This demonstrates the Mcm2 antibody detected the Mcm2 protein (arrow).

Mouse Mcm2 Antibody (C-term) - Background

Mcm2 acts as a factor that allows the DNA to undergo a single round of replication per cell cycle. Required for the entry in S phase and for cell division.

Mouse Mcm2 Antibody (C-term) - References

- Kunnev, D., et al. Oncogene 29(25):3630-3638(2010)
- Chuang, C.H., et al. PLoS Genet. 6 (9) (2010) :
- Lee, E.Y., et al. Dev. Biol. 332(1):104-115(2009)
- Hasegawa, M., et al. Leuk. Res. 33(8):1100-1107(2009)
- Breunig, J.J., et al. Proc. Natl. Acad. Sci. U.S.A. 105(35):13127-13132(2008)

Precautions

Mouse Mcm2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Mcm2 Antibody (C-term) - Protein Information

Name Mcm2

Synonyms Bm28, Cdcl1, Kiaa0030, Mcmd2

Function

Acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity. Required for the entry in S phase and for cell division. Plays a role in terminally differentiated hair cells development of the cochlea and induces cells apoptosis.

Cellular Location

Nucleus {ECO:0000250|UniProtKB:P49736}.

Chromosome

{ECO:0000250|UniProtKB:P55861}.

Note=Associated with chromatin before the formation of nuclei and detaches from it as DNA replication progresses.

{ECO:0000250|UniProtKB:P55861}

Mouse Mcm2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)