

ULK1 Antibody (Center S317)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19250c

Specification

ULK1 Antibody (Center S317) - Product Information

Application	WB,E
Primary Accession	O75385
Other Accession	NP_003556.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	296-322

ULK1 Antibody (Center S317) - Additional Information

Gene ID 8408

Other Names

Serine/threonine-protein kinase ULK1,
Autophagy-related protein 1 homolog,
ATG1, hATG1, Unc-51-like kinase 1, ULK1,
KIAA0722

Target/Specificity

This ULK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 296-322 amino acids from the Central region of human ULK1.

Dilution

WB~~1:2000


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



All lanes : Anti-ULK1 at 1:2000 dilution
Lane 1: HeLa whole cell lysate
Lane 2: RD whole cell lysate
Lane 3: C2C12 whole cell lysate
Lane 4: NIH/3T3 whole cell lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L),
Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 113 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

ULK1 Antibody (Center S317) - Background

ULK1 is involved in autophagy. Required for autophagosome formation (By similarity). Target of the TOR kinase signaling pathway that regulates autophagy through the control of phosphorylation status of ATG13/KIAA0652 and ULK1, and the regulation of the ATG13-ULK1-RB1CC1 complex (By similarity). Phosphorylates ATG13/KIAA0652. Involved in axon growth (By similarity). Plays an essential role in neurite extension of cerebellar granule cells (By similarity).

ULK1 Antibody (Center S317) - References

Mercer, C.A., et al. Autophagy

ULK1 Antibody (Center S317) is for research use only and not for use in diagnostic or therapeutic procedures.

ULK1 Antibody (Center S317) - Protein Information

Name ULK1 ([HGNC:12558](#))

Synonyms KIAA0722

Function

Serine/threonine-protein kinase involved in autophagy in response to starvation (PubMed:18936157, PubMed:21460634, PubMed:21795849, PubMed:25040165). Acts upstream of phosphatidylinositol 3-kinase PIK3C3 to regulate the formation of autophagophores, the precursors of autophagosomes (PubMed:18936157, PubMed:21460634, PubMed:21795849, PubMed:25040165). Part of regulatory feedback loops in autophagy: acts both as a downstream effector and negative regulator of mammalian target of rapamycin complex 1 (mTORC1) via interaction with RPTOR (PubMed:21795849). Activated via phosphorylation by AMPK and also acts as a regulator of AMPK by mediating phosphorylation of AMPK subunits PRKAA1, PRKAB2 and PRKAG1, leading to negatively regulate AMPK activity (PubMed:21460634). May

5(5):649-662(2009)
Ganley, I.G., et al. J. Biol. Chem. 284(18):12297-12305(2009)
Jung, C.H., et al. Mol. Biol. Cell 20(7):1992-2003(2009)
Hosokawa, N., et al. Mol. Biol. Cell 20(7):1981-1991(2009)
Chan, E.Y. Sci Signal 2 (84), PE51 (2009) :

phosphorylate ATG13/KIAA0652 and RPTOR; however such data need additional evidences (PubMed:18936157). Plays a role early in neuronal differentiation and is required for granule cell axon formation (PubMed:11146101). May also phosphorylate SESN2 and SQSTM1 to regulate autophagy (PubMed:25040165). Phosphorylates FLCN, promoting autophagy (PubMed:25126726).

Cellular Location

Cytoplasm, cytosol. Preautophagosomal structure. Note=Under starvation conditions, is localized to punctate structures primarily representing the isolation membrane that sequesters a portion of the cytoplasm resulting in the formation of an autophagosome.

Tissue Location

Ubiquitously expressed. Detected in the following adult tissues: skeletal muscle, heart, pancreas, brain, placenta, liver, kidney, and lung

ULK1 Antibody (Center S317) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ULK1 Antibody (Center S317) - Citations

- [Neratinib decreases pro-survival responses of \[sorafenib + vorinostat\] in pancreatic cancer](#)