

**ENAH Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP19271B**

### Specification

#### ENAH Antibody (C-term) - Product Information

Application	WB, E
Primary Accession	<a href="#">Q8N8S7</a>
Other Accession	<a href="#">NP_060682.2</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	66510
Antigen Region	446-475

#### ENAH Antibody (C-term) - Additional Information

##### Gene ID 55740

##### Other Names

Protein enabled homolog, ENAH, MENA

##### Target/Specificity

This ENAH antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 446-475 amino acids from the C-terminal region of human ENAH.

##### Dilution

WB~1:1000

##### Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

##### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

ENAH Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.



ENAH Antibody (C-term)(Cat. #AP19271b) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the ENAH antibody detected the ENAH protein (arrow).

#### ENAH Antibody (C-term) - Background

Ena/VASP proteins are actin-associated proteins involved in a range of processes dependent on cytoskeleton remodeling and cell polarity such as axon guidance and lamellipodial and filopodial dynamics in migrating cells. ENAH induces the formation of F-actin rich outgrowths in fibroblasts. Acts synergistically with BAIAP2-alpha and downstream of NTN1 to promote filopodia formation. Required for the actin-based mobility of Listeria monocytogenes (By similarity).

#### ENAH Antibody (C-term) - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :  
Chattopadhyay, I., et al. Mutat. Res. 696(2):130-138(2010)  
Hahn, W.H., et al. Exp. Mol. Med. 41(11):793-801(2009)  
Gurzu, S., et al. Rom J Morphol Embryol 50(2):213-216(2009)

**ENAH Antibody (C-term) - Protein Information**

Higashi, M., et al. PLoS ONE 4 (3), E4765 (2009) :

**Name** ENAH

**Synonyms** MENA

**Function**

Ena/VASP proteins are actin-associated proteins involved in a range of processes dependent on cytoskeleton remodeling and cell polarity such as axon guidance and lamellipodial and filopodial dynamics in migrating cells. ENAH induces the formation of F-actin rich outgrowths in fibroblasts. Acts synergistically with BAIAP2-alpha and downstream of NTN1 to promote filopodia formation (By similarity).

**Cellular Location**

Cytoplasm. Cytoplasm, cytoskeleton. Cell projection, lamellipodium. Cell projection, filopodium. Cell junction, synapse. Cell junction, focal adhesion. Note=Targeted to the leading edge of lamellipodia and filopodia by MRL family members. Colocalizes at filopodial tips with a number of other proteins including vinculin and zyxin. Colocalizes with N-WASP at the leading edge. Colocalizes with GPHN and PFN at synapses (By similarity).

**Tissue Location**

Expressed in myoepithelia of parotid, breast, bronchial glands and sweat glands. Expressed in colon-rectum muscularis mucosae epithelium, pancreas acinar ductal epithelium, endometrium epithelium, prostate fibromuscular stroma and placenta vascular media Overexpressed in a majority of breast cancer cell lines and primary breast tumor lesions.

**ENAH Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)