

SQSTM1 Antibody(C-term S403)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19505b

Specification

SQSTM1 Antibody(C-term S403) - Product Information

Application	WB,E
Primary Accession	Q13501
Other Accession	Q08623 , Q64337 , NP_001135770.1
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	47687
Antigen Region	383-409

SQSTM1 Antibody(C-term S403) - Additional Information

Gene ID 8878

Other Names

Sequestosome-1, EBI3-associated protein of 60 kDa, EBIAP, p60, Phosphotyrosine-independent ligand for the Lck SH2 domain of 62 kDa, Ubiquitin-binding protein p62, SQSTM1, ORCA, OSIL

Target/Specificity

This SQSTM1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 383-409 amino acids from the C-terminal region of human SQSTM1.

Dilution


WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2



SQSTM1 Antibody (C-term S403) (Cat. #AP19505b) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the SQSTM1 antibody detected the SQSTM1 protein (arrow).

SQSTM1 Antibody(C-term S403) - Background

This gene encodes a multifunctional protein that binds ubiquitin and regulates activation of the nuclear factor kappa-B (NF-κB) signaling pathway. The protein functions as a scaffolding/adaptor protein in concert with TNF receptor-associated factor 6 to mediate activation of NF-κB in response to upstream signals. Alternatively spliced transcript variants encoding either the same or different isoforms have been identified for this gene. Mutations in this gene result in sporadic and familial Paget disease of bone.

SQSTM1 Antibody(C-term S403) - References

Visconti, M.R., et al. J. Bone Miner. Res.

weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SQSTM1 Antibody(C-term S403) is for research use only and not for use in diagnostic or therapeutic procedures.

25(11):2368-2373(2010)
Ding, W.X., et al. J. Biol. Chem.
285(36):27879-27890(2010)
Gao, C., et al. Nat. Cell Biol.
12(8):781-790(2010)
Jain, A., et al. J. Biol. Chem.
285(29):22576-22591(2010)
Lau, A., et al. Mol. Cell. Biol.
30(13):3275-3285(2010)

SQSTM1 Antibody(C-term S403) - Protein Information

Name SQSTM1

Synonyms ORCA, OSIL

Function

Autophagy receptor required for selective macroautophagy (aggrephagy). Functions as a bridge between polyubiquitinated cargo and autophagosomes. Interacts directly with both the cargo to become degraded and an autophagy modifier of the MAP1 LC3 family (PubMed:16286508, PubMed:20168092, PubMed:24128730, PubMed:28404643, PubMed:22622177). Along with WDFY3, involved in the formation and autophagic degradation of cytoplasmic ubiquitin-containing inclusions (p62 bodies, ALIS/aggresome-like induced structures). Along with WDFY3, required to recruit ubiquitinated proteins to PML bodies in the nucleus (PubMed:24128730, PubMed:20168092). May regulate the activation of NFKB1 by TNF-alpha, nerve growth factor (NGF) and interleukin-1. May play a role in titin/TTN downstream signaling in muscle cells. May regulate signaling cascades through ubiquitination. Adapter that mediates the

interaction between TRAF6 and CYLD (By similarity). May be involved in cell differentiation, apoptosis, immune response and regulation of K(+) channels. Involved in endosome organization by retaining vesicles in the perinuclear cloud: following ubiquitination by RNF26, attracts specific vesicle-associated adapters, forming a molecular bridge that restrains cognate vesicles in the perinuclear region and organizes the endosomal pathway for efficient cargo transport (PubMed:27368102). Promotes relocalization of 'Lys-63'-linked ubiquitinated STING1 to autophagosomes (PubMed:29496741). Acts as an activator of the NFE2L2/NRF2 pathway via interaction with KEAP1: interaction inactivates the BCR(KEAP1) complex, promoting nuclear accumulation of NFE2L2/NRF2 and subsequent expression of cytoprotective genes (PubMed:20452972, PubMed:28380357).

Cellular Location

Cytoplasm, cytosol. Late endosome. Lysosome. Cytoplasmic vesicle, autophagosome. Nucleus. Endoplasmic reticulum. Nucleus, PML body. Cytoplasm, myofibril, sarcomere. Note=In cardiac muscle, localizes to the sarcomeric band (By similarity). Commonly found in inclusion bodies containing polyubiquitinated protein aggregates. In neurodegenerative diseases, detected in Lewy bodies in Parkinson disease, neurofibrillary tangles in Alzheimer disease, and HTT aggregates in Huntington disease. In protein aggregate diseases of the liver, found in large amounts in Mallory bodies of alcoholic and nonalcoholic steatohepatitis, hyaline bodies in hepatocellular carcinoma, and in SERPINA1 aggregates Enriched in Rosenthal fibers of pilocytic astrocytoma. In the cytoplasm, observed in both membrane-free ubiquitin-containing protein aggregates (sequestosomes) and membrane-surrounded autophagosomes Colocalizes with TRIM13 in the perinuclear endoplasmic reticulum. Co- localizes with

TRIM5 in cytoplasmic bodies. When nuclear export is blocked by treatment with leptomycin B, accumulates in PML bodies

Tissue Location

Ubiquitously expressed.

**SQSTM1 Antibody(C-term S403) -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)