

EIF4E2 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1955a

Specification

EIF4E2 Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O60573
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	28362
Antigen Region	12-41

EIF4E2 Antibody (N-term) - Additional Information

Gene ID 9470

Other Names

Eukaryotic translation initiation factor 4E type 2, eIF-4E type 2, eIF4E type 2, Eukaryotic translation initiation factor 4E homologous protein, Eukaryotic translation initiation factor 4E-like 3, eIF4E-like protein 4E-LP, mRNA cap-binding protein 4EHP, mRNA cap-binding protein type 3, EIF4E2, EIF4EL3

Target/Specificity

This EIF4E2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 12-41 amino acids from the N-terminal region of human EIF4E2.

Dilution


WB~~1:1000
IHC-P~~1:50~100

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C



Western blot analysis of EIF4E2 Antibody (N-term) polyclonal antibody (Cat.#AP1955a) (arrow). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the EIF4E2 gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human breast carcinoma reacted with EIF4E2 Antibody (N-term) (Cat.#AP1955a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

EIF4E2 Antibody (N-term) - Background

in small aliquots to prevent freeze-thaw cycles.

Precautions

EIF4E2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

EIF4E2 Antibody (N-term) - Protein Information

Name EIF4E2

{ECO:0000303|PubMed:15153109,
ECO:0000312|HGNC:HGNC:3293}

Function

Recognizes and binds the 7-methylguanosine-containing mRNA cap during an early step in the initiation (PubMed:17368478, PubMed:25624349, PubMed:9582349). Acts as a repressor of translation initiation (PubMed:22751931). In contrast to EIF4E, it is unable to bind eIF4G (EIF4G1, EIF4G2 or EIF4G3), suggesting that it acts by competing with EIF4E and block assembly of eIF4F at the cap (By similarity). In P-bodies, component of a complex that promotes miRNA- mediated translational repression (PubMed:28487484).

Cellular Location

Cytoplasm. Cytoplasm, P-body

EIF4E2 is expressed exclusively in the cytoplasm. This protein recognizes and binds the 7 methylguanosine containing mRNA cap during an early step in the initiation of protein synthesis and facilitates ribosome binding by inducing the unwinding of the mRNAs secondary structures.

EIF4E2 Antibody (N-term) - References

Rom, E., et al., J. Biol. Chem. 273(21):13104-13109 (1998).
Mao, M., et al., Proc. Natl. Acad. Sci. U.S.A. 95(14):8175-8180 (1998).
Tee, A.R., et al., FEBS Lett. 564 (1-2), 58-62 (2004) ().

EIF4E2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)