

## **BCL2 Antibody(Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP19560c

## **Specification**

#### **BCL2 Antibody(Center) - Product Information**

Application WB,E
Primary Accession P10415

Other Accession P49950, P10417,

<u>0911V8</u>, <u>000709</u>,

O02718, NP\_000624.2

Reactivity Human

Predicted Bovine, Chicken,

Hamster, Mouse,

Rat

Host Rabbit
Clonality Polyclonal
Isotype Rabbit Ig
Calculated MW 26266
Antigen Region 144-171

**BCL2 Antibody(Center) - Additional Information** 

## Gene ID 596

# **Other Names**

Apoptosis regulator Bcl-2, BCL2

### **Target/Specificity**

This BCL2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 144-171 amino acids from the Central region of human BCL2.

## **Dilution**


WB~~1:1000

### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### **Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



BCL2 Antibody (Center) (Cat. #AP19560c) western blot analysis in MDA-MB453 cell line lysates (35ug/lane). This demonstrates the BCL2 antibody detected the BCL2 protein (arrow).

## BCL2 Antibody(Center) - Background

This gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Two transcript variants, produced by alternate splicing, differ in their C-terminal ends. [provided by RefSeq].

## **BCL2 Antibody(Center) - References**

Feng, H., et al. Cancer Cell 18(4):353-366(2010)

Azad, N., et al. Ann. N. Y. Acad. Sci. 1203, 1-6 (2010):

Dubikov, A.I., et al. Scand. J. Rheumatol.

39(5):368-372(2010)

Yu, B., et al. J. Exp. Clin. Cancer Res. 29, 107





### **Precautions**

BCL2 Antibody(Center) is for research use only and not for use in diagnostic or therapeutic procedures.

(2010):

Trisciuoglio, D., et al. PLoS ONE 5 (7), E11772 (2010) :

## **BCL2 Antibody(Center) - Protein Information**

## Name BCL2

#### **Function**

Suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural cells. Regulates cell death by controlling the mitochondrial membrane permeability. Appears to function in a feedback loop system with caspases. Inhibits caspase activity either by preventing the release of cytochrome c from the mitochondria and/or by binding to the apoptosis-activating factor (APAF-1). May attenuate inflammation by impairing NLRP1-inflammasome activation, hence CASP1 activation and IL1B release (PubMed:<a href="http://www.uniprot.org/c itations/17418785" target="\_blank">17418785</a>).

## **Cellular Location**

Mitochondrion outer membrane; Single-pass membrane protein. Nucleus membrane; Single-pass membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein

# **Tissue Location**

Expressed in a variety of tissues.

## **BCL2 Antibody(Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture