

DNAJB12 Antibody(N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19585a

Specification

DNAJB12 Antibody(N-term) - Product Information

Application	WB, E
Primary Accession	Q9NXW2
Other Accession	Q58DR2
Reactivity	Human
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	41860
Antigen Region	7-36

DNAJB12 Antibody(N-term) - Additional Information

Gene ID 54788

Other Names

Dnaj homolog subfamily B member 12,
 DNAJB12

Target/Specificity

This DNAJB12 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 7-36 amino acids from the N-terminal region of human DNAJB12.

Dilution

WB~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DNAJB12 Antibody(N-term) is for research



DNAJB12 Antibody (N-term) (Cat. #AP19585a) western blot analysis in T47D cell line lysates (35ug/lane). This demonstrates the DNAJB12 antibody detected the DNAJB12 protein (arrow).

DNAJB12 Antibody(N-term) - Background

DNAJB12 belongs to the evolutionarily conserved DNAJ/HSP40 family of proteins, which regulate molecular chaperone activity by stimulating ATPase activity. DNAJ proteins may have up to 3 distinct domains: a conserved 70-amino acid J domain, usually at the N terminus; a glycine/phenylalanine (G/F)-rich region; and a cysteine-rich domain containing 4 motifs resembling a zinc finger domain (Ohtsuka and Hata, 2000 [PubMed 11147971]). [supplied by OMIM].

DNAJB12 Antibody(N-term) - References

Lamesch, P., et al. Genomics 89(3):307-315(2007)

Ohtsuka, K., et al. Cell Stress Chaperones 5(2):98-112(2000)

use only and not for use in diagnostic or therapeutic procedures.

DNAJB12 Antibody(N-term) - Protein Information

Name DNAJB12

{ECO:0000303|PubMed:21150129,
ECO:0000312|HGNC:HGNC:14891}

Function

Acts as a co-chaperone with HSPA8/Hsc70; required to promote protein folding and trafficking, prevent aggregation of client proteins, and promote unfolded proteins to endoplasmic reticulum- associated degradation (ERAD) pathway (PubMed:21150129, PubMed:21148293). Acts by determining HSPA8/Hsc70's ATPase and polypeptide-binding activities (PubMed:21148293). Can also act independently of HSPA8/Hsc70: together with DNAJB14, acts as a chaperone that promotes maturation of potassium channels KCND2 and KCNH2 by stabilizing nascent channel subunits and assembling them into tetramers (PubMed:27916661). While stabilization of nascent channel proteins is dependent on HSPA8/Hsc70, the process of oligomerization of channel subunits is independent of HSPA8/Hsc70 (PubMed:27916661). When overexpressed, forms membranous structures together with DNAJB14 and HSPA8/Hsc70 within the nucleus; the role of these structures, named DJANGOs, is still unclear (PubMed:24732912).

Cellular Location

Endoplasmic reticulum membrane;
Single-pass membrane protein. Nucleus membrane; Single-pass membrane protein.
Note=Localizes to the endoplasmic reticulum membrane (PubMed:21150129, PubMed:21148293, PubMed:24732912, PubMed:27916661) When overexpressed, forms membranous structures in the

nucleus (PubMed:24732912).

DNAJB12 Antibody(N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)