

**NFKB1 Antibody (S932)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP1980c**

### Specification

#### NFKB1 Antibody (S932) - Product Information

Application	WB, E
Primary Accession	<a href="#">P19838</a>
Other Accession	<a href="#">P25799</a> , <a href="#">Q04861</a>
Reactivity	Human
Predicted	Chicken, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	105356
Antigen Region	911-939

#### NFKB1 Antibody (S932) - Additional Information

Gene ID 4790

#### Other Names

Nuclear factor NF-kappa-B p105 subunit, DNA-binding factor KBF1, EBP-1, Nuclear factor of kappa light polypeptide gene enhancer in B-cells 1, Nuclear factor NF-kappa-B p50 subunit, NFKB1

#### Target/Specificity

This NFKB1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 911-939 amino acids from human NFKB1.

#### Dilution

WB~1:1000


#### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions



All lanes : Anti-NFKB1 Antibody (S932) at 1:1000 dilution Lane 1: Raji whole cell lysate  
Lane 2: A431 whole cell lysate  
Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L),  
Peroxidase conjugated at 1/10000 dilution.  
Predicted band size : 105 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

#### NFKB1 Antibody (S932) - Background

NFKB1 is a 105 kD protein which can undergo cotranslational processing by the 26S proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFKB) protein complex. NFKB is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NFKB translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NFKB has been associated with a number of inflammatory diseases while persistent

NFKB1 Antibody (S932) is for research use only and not for use in diagnostic or therapeutic procedures.

#### NFKB1 Antibody (S932) - Protein Information

##### Name NFKB1

##### Function

NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain- containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I- kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. NF-kappa-B heterodimeric p65-p50 and RelB-p50 complexes are transcriptional activators. The NF-kappa-B p50-p50 homodimer is a transcriptional repressor, but can act as a transcriptional activator when associated with BCL3. NFKB1 appears to have dual functions such as cytoplasmic retention of attached NF-kappa-B proteins by p105 and generation of p50 by a cotranslational processing. The proteasome-mediated process ensures the production of both p50

inhibition of NFkB leads to inappropriate immune cell development or delayed cell growth.

#### NFKB1 Antibody (S932) - References

Legembre, P., et al., *J. Biol. Chem.* 279(45):46742-46747 (2004).  
Imamura, R., et al., *J. Biol. Chem.* 279(45):46415-46423 (2004).  
Hung, J.H., et al., *J. Biol. Chem.* 279(45):46384-46392 (2004).  
Binnicker, M.J., et al., *Infect. Immun.* 72(11):6408-6417 (2004).  
Zhang, H., et al., *J. Biol. Chem.* 279(43):44955-44965 (2004).

and p105 and preserves their independent function, although processing of NFKB1/p105 also appears to occur post-translationally. p50 binds to the kappa-B consensus sequence 5'-GGRNNYYCC-3', located in the enhancer region of genes involved in immune response and acute phase reactions. In a complex with MAP3K8, NFKB1/p105 represses MAP3K8-induced MAPK signaling; active MAP3K8 is released by proteasome-dependent degradation of NFKB1/p105.

#### **Cellular Location**

Nucleus. Cytoplasm. Note=Nuclear, but also found in the cytoplasm in an inactive form complexed to an inhibitor (I- kappa-B)

#### **NFKB1 Antibody (S932) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)