

CD73 (NT5E) Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2014A

Specification

CD73 (NT5E) Antibody (N-term) - Product Information

Application	IHC-P,E
Primary Accession	P21589
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	1-30

CD73 (NT5E) Antibody (N-term) - Additional Information

Gene ID 4907

Other Names

5'-nucleotidase, 5'-NT, Ecto-5'-nucleotidase, CD73, NT5E, NT5, NTE

Target/Specificity

This CD73 (NT5E) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human CD73 (NT5E).

Dilution

IHC-P~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CD73 (NT5E) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.



CD73 (NT5E) Antibody (N-term) (Cat. #AP2014a) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CD73 (NT5E) Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

CD73 (NT5E) Antibody (N-term) - Background

Ecto-5-prime-nucleotidase catalyzes conversion at neutral pH of purine 5-prime mononucleotides to nucleosides. The enzyme consists of a dimer of 2 identical 70 kD subunits tethered by a glycosyl phosphatidyl inositol moiety to the exterior plasma membrane surface. The enzyme, a marker of lymphocyte differentiation is associated in deficiency with a variety of immunodeficiency diseases. The preferred substrate is AMP. The NT5 gene has been localized to 6q14-q21 by immunofluorescence and a study of a panel of human x mouse hybrids that contained fragments of chromosome 6 as translocations.

CD73 (NT5E) Antibody (N-term) - Protein Information**Name** NT5E**Synonyms** NT5, NTE**Function**

Hydrolyzes extracellular nucleotides into membrane permeable nucleosides. Exhibits AMP-, NAD-, and NMN-nucleosidase activities.

Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor.

CD73 (NT5E) Antibody (N-term) - References

Hashikawa, T., et al., J. Dent. Res. 82(11):888-892 (2003).
Rosi, F., et al., Biomed. Pharmacother. 56(2):100-104 (2002).
Misumi, Y., et al., Eur. J. Biochem. 191(3):563-569 (1990).
Boyle, J.M., et al., Hum. Genet. 81(1):88-92 (1988).
Kalsi, K., et al., Mol. Cell. Biochem. 232 (1-2), 113-119 (2002).

CD73 (NT5E) Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD73 (NT5E) Antibody (N-term) - Citations

- [Umbilical cord mesenchymal stromal cells affected by gestational diabetes mellitus display premature aging and mitochondrial dysfunction.](#)
- [CD73 is a phenotypic marker of effector memory Th17 cells in inflammatory bowel disease.](#)
- [Ecto-5'-nucleotidase/CD73 knockdown increases cell migration and mRNA level of collagen I in a hepatic stellate cell line.](#)
- [Expression of dopamine-associated genes on conjunctiva stromal-derived human mesenchymal stem cells.](#)