

**PROX1 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2035C**

**Specification**

**PROX1 Antibody (Center) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">Q92786</a>
Other Accession	<a href="#">P48437</a> , <a href="#">NP_002754</a>
Reactivity	Human, Mouse
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	185-214

**PROX1 Antibody (Center) - Additional Information**

**Gene ID** 5629

**Other Names**

Prospero homeobox protein 1, Homeobox prospero-like protein PROX1, PROX-1, PROX1

**Target/Specificity**

This PROX1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 185-214 amino acids from the Central region of human PROX1.

**Dilution**

WB~~1:500  
 IHC-P~~1:10~50

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PROX1 Antibody (Center) is for research use



All lanes : Anti-PROX1 Antibody (Center) at 1:500 dilution  
 Lane 1: HepG2 whole cell lysate  
 Lane 2: SH-SY5Y, whole cell lysate  
 Lane 3: SW620 whole cell lysate  
 Lysates/proteins at 20 µg per lane.  
 Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.  
 Predicted band size : 83 kDa  
 Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-PROX1 Antibody (Center) at 1:500 dilution + Mouse heart whole cell lysate

only and not for use in diagnostic or therapeutic procedures.

#### PROX1 Antibody (Center) - Protein Information

**Name** PROX1

#### Function

Transcription factor involved in developmental processes such as cell fate determination, gene transcriptional regulation and progenitor cell regulation in a number of organs. Plays a critical role in embryonic development and functions as a key regulatory protein in neurogenesis and the development of the heart, eye lens, liver, pancreas and the lymphatic system. Involved in the regulation of the circadian rhythm. Represses: transcription of the retinoid-related orphan receptor RORG, transcriptional activator activity of RORA and RORG and the expression of RORA/G-target genes including core clock components: ARNTL/BMAL1, NPAS2 and CRY1 and metabolic genes: AVPR1A and ELOVL3.

#### Cellular Location

Nucleus {ECO:0000250|UniProtKB:P48437}.  
Note=RORG promotes its nuclear localization.  
{ECO:0000250|UniProtKB:P48437}

#### Tissue Location


Most actively expressed in the developing lens. Detected also in embryonic brain, lung, liver and kidney. In adult, it is more abundant in heart and liver than in brain, skeletal muscle, kidney and pancreas.

#### PROX1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L),  
Peroxidase conjugated at 1/10000 dilution.  
Predicted band size : 83 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with PROX1 antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

#### PROX1 Antibody (Center) - Background

The expression pattern of Prox1 suggests that it has a role in a variety of embryonic tissues, including lens. Prox mRNA is present in many different human tissues with lens demonstrating the highest level. Homozygous Prox1-null mice die at midgestation from multiple developmental defects, and a targeted effect on lens development has been reported. Prox1 inactivation caused abnormal cellular proliferation, downregulated expression of the cell cycle inhibitors Cdkn1b and Cdkn1c, misexpression of E-cadherin, and excessive apoptosis. Consequently, mutant lens cells failed to polarize and elongate properly, resulting in a hollow lens. Prox1 is expressed in a subpopulation of endothelial cells that by budding and sprouting give rise to the lymphatic system. Prox1 appears to be a specific and required regulator of the development of the lymphatic system. Prox1 also has been documented to be required for hepatocyte migration in the mouse. Loss of Prox1 results in a smaller liver with a reduced

population of clustered hepatocytes. The homeodomain protein Prox1 regulates the egress of progenitor cells from the cell cycle in the embryonic mouse retina. Cells lacking Prox1 are less likely to stop dividing, and ectopic expression of Prox1 forces progenitor cells to exit the cell cycle. Prox1 acts as a key participant in progenitor-cell proliferation and cell-fate determination in the vertebrate retina.

#### **PROX1 Antibody (Center) - References**

Nagai, H., et al., *Genes Chromosomes Cancer* 38(1):13-21 (2003).

Dyer, M.A., et al., *Nat. Genet.* 34(1):53-58 (2003).

Hong, Y.K., et al., *Dev. Dyn.* 225(3):351-357 (2002).

Petrova, T.V., et al., *EMBO J.* 21(17):4593-4599 (2002).

Mouta Carreira, C., et al., *Cancer Res.* 61(22):8079-8084 (2001).