

P100 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2044a

Specification

P100 Antibody - Product Information

Application	WB, IHC-P,E
Primary Accession	Q9ULW0
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	85653
Antigen Region	350-382

P100 Antibody - Additional Information

Gene ID 22974

Other Names

Targeting protein for Xklp2, Differentially expressed in cancerous and non-cancerous lung cells 2, DIL-2, Hepatocellular carcinoma-associated antigen 519, Hepatocellular carcinoma-associated antigen 90, Protein fls353, Restricted expression proliferation-associated protein 100, p100, TPX2, C20orf1, C20orf2, DIL2, HCA519

Target/Specificity

This P100 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 350-382 amino acids from human P100.

Dilution


WB~~1:1000
IHC-P~~1:50~100

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw



The anti-p100 Pab (Cat. #AP2044a) is used in Western blot to detect p100 in placenta lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

P100 Antibody - Background

The nuclear protein p100 is a proliferation-associated protein whose

cycles.

Precautions

P100 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

P100 Antibody - Protein Information

Name TPX2

Synonyms C20orf1, C20orf2, DIL2, HCA519

Function

Spindle assembly factor required for normal assembly of mitotic spindles. Required for normal assembly of microtubules during apoptosis. Required for chromatin and/or kinetochore dependent microtubule nucleation. Mediates AURKA localization to spindle microtubules (PubMed:18663142, PubMed:19208764). Activates AURKA by promoting its autophosphorylation at 'Thr-288' and protects this residue against dephosphorylation (PubMed:18663142, PubMed:19208764). TPX2 is inactivated upon binding to importin-alpha (PubMed:26165940). At the onset of mitosis, GOLGA2 interacts with importin-alpha, liberating TPX2 from importin-alpha, allowing TPX2 to activate AURKA kinase and stimulates local microtubule nucleation (PubMed:26165940).

Cellular Location

Nucleus. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, spindle pole. Note=During mitosis it is strictly associated with the spindle pole and with the mitotic spindle, whereas during S and G2, it is diffusely distributed throughout the nucleus. Is released from the nucleus in apoptotic cells and is detected on apoptotic

expression is restricted to cell cycle phases S, G2, and M. Exclusively expressed in proliferating cells from the transition G1/S until the end of cytokinesis. During mitosis it is strictly associated with the spindle pole and with the mitotic spindle, whereas during S and G2, it is diffusely distributed throughout the nucleus. The full-length cDNA encodes a 747-amino acid protein with a putative ATP/GTP-binding site motif. RT-PCR analysis demonstrated strong expression of in lung carcinoma cell lines but not in normal lung tissues. Expression was also found in adult placenta, skeletal muscle, thymus, testis, and small intestine and in fetal brain, liver, and kidney. P100 is also correlated to cancer prognosis.

P100 Antibody - References

Ota, T., et al., Nat. Genet. 36(1):40-45 (2004).
Heidebrecht, H.J., et al., Mol. Cancer Res. 1(4):271-279 (2003).
Garrett, S., et al., Curr. Biol. 12(23):2055-2059 (2002).
Gruss, O.J., et al., Nat. Cell Biol. 4(11):871-879 (2002).
Kufer, T.A., et al., J. Cell Biol. 158(4):617-623 (2002).

microtubules

Tissue Location

Expressed in lung carcinoma cell lines but
not in normal lung tissues

P100 Antibody - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)