

CDKN1A Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20643c

Specification

CDKN1A Antibody (C-term) - Product Information

Application	IF, WB, IHC-P,E
Primary Accession	P38936
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig

CDKN1A Antibody (C-term) - Additional Information

Gene ID 1026

Other Names

Cyclin-dependent kinase inhibitor 1,
CDK-interacting protein 1, Melanoma
differentiation-associated protein 6, MDA-6,
p21, CDKN1A, CAP20, CDKN1, CIP1, MDA6,
PIC1, SDI1, WAF1

Target/Specificity

This CDKN1A antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 133-164 amino acids from the C-terminal region of human CDKN1A.

Dilution

IF~~1:25
WB~~1:1000
IHC-P~~1:25


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



Fluorescent image of A549 cells stained with CDKN1A Antibody (C-term)(Cat#AP20643c). AP20643c was diluted at 1:25 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red). Cytoplasm



All lanes : Anti-CDKN1A Antibody (C-term) at 1:2000 dilution Lane 1: 293 whole cell lysate

CDKN1A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CDKN1A Antibody (C-term) - Protein Information

Name CDKN1A

Synonyms CAP20, CDKN1, CIP1, MDA6, PIC1, SDI1, WA

Function

May be involved in p53/TP53 mediated inhibition of cellular proliferation in response to DNA damage. Binds to and inhibits cyclin- dependent kinase activity, preventing phosphorylation of critical cyclin-dependent kinase substrates and blocking cell cycle progression. Functions in the nuclear localization and assembly of cyclin D-CDK4 complex and promotes its kinase activity towards RB1. At higher stoichiometric ratios, inhibits the kinase activity of the cyclin D- CDK4 complex. Inhibits DNA synthesis by DNA polymerase delta by competing with POLD3 for PCNA binding (PubMed:11595739). Plays an important role in controlling cell cycle progression and DNA damage- induced G2 arrest (PubMed:9106657).

Cellular Location

Cytoplasm. Nucleus

Tissue Location


Expressed in all adult tissues, with 5-fold lower levels observed in the brain

CDKN1A Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

Lane 2: HepG2 whole cell lysate Lane 3: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 18 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Western blot analysis of lysate from HeLa cell line, using CDKN1A Antibody (C-term)(Cat. #AP20797c). AP20797c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 35ug.



Western blot analysis of lysates from HeLa, HUVEC, MCF-7, SH-SY5Y cell line and mouse lung tissue lysate(from left to right), using CDKN1A Antibody (C-term)(Cat. #AP20643c). AP20643c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

- [Flow Cytometry](#)
- [Cell Culture](#)



AP20643C staining CDKN1A in human colon tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

CDKN1A Antibody (C-term) - Background

May be the important intermediate by which p53/TP53 mediates its role as an inhibitor of cellular proliferation in response to DNA damage. Binds to and inhibits cyclin-dependent kinase activity, preventing phosphorylation of critical cyclin-dependent kinase substrates and blocking cell cycle progression. Functions in the nuclear localization and assembly of cyclin D- CDK4 complex and promotes its kinase activity towards RB1. At higher stoichiometric ratios, inhibits the kinase activity of the cyclin D-CDK4 complex.

CDKN1A Antibody (C-term) - References

Harper J.W., et al. Cell 75:805-816(1993).
El-Deiry W.S., et al. Cell 75:817-825(1993).
Xiong Y., et al. Nature 366:701-704(1993).
Jiang H., et al. Mol. Cell. Differ. 1:285-299(1993).
Jiang H., et al. Oncogene 10:1855-1864(1995).