

**BMP9 (GDF2) Antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2064A**

**Specification**

**BMP9 (GDF2) Antibody (N-term) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">Q9UK05</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	8-37

**BMP9 (GDF2) Antibody (N-term) - Additional Information**

Gene ID 2658

**Other Names**

Growth/differentiation factor 2, GDF-2, Bone morphogenetic protein 9, BMP-9, GDF2, BMP9

**Target/Specificity**

This BMP9 (GDF2) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 8-37 amino acids from the N-terminal region of human BMP9 (GDF2).

**Dilution**

WB~~1:1000  
 IHC-P~~1:50~100

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

BMP9 (GDF2) Antibody (N-term) is for



The anti-hGDF2 N-term Pab (Cat. #AP2064a) is used in Western blot to detect hGDF2 in mouse liver tissue lysate.



Western blot analysis of GDF2 (arrow) using rabbit polyclonal GDF2-K23 (Cat. #AP2064a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the GDF2 gene.

research use only and not for use in diagnostic or therapeutic procedures.

#### BMP9 (GDF2) Antibody (N-term) - Protein Information

**Name** GDF2

**Synonyms** BMP9

#### Function


Potent circulating inhibitor of angiogenesis. Signals through the type I activin receptor ACVRL1 but not other Alks. Signaling through SMAD1 in endothelial cells requires TGF-beta coreceptor endoglin/ENG.

#### Cellular Location

Secreted

#### Tissue Location

Detected in blood plasma (at protein level).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

#### BMP9 (GDF2) Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### BMP9 (GDF2) Antibody (N-term) - Background

GDF2 is a member of the bone morphogenetic protein (BMP) family and the TGF-beta superfamily. This group of proteins is characterized by a polybasic proteolytic processing site which is cleaved to produce a mature protein containing seven conserved cysteine residues. The members of this family are regulators of cell growth and differentiation in both embryonic and adult tissues. Studies in rodents suggest that this protein plays a role in the adult liver and in differentiation of cholinergic central nervous system neurons.

#### BMP9 (GDF2) Antibody (N-term) - References

Majumdar, M.K., et al., *J. Cell. Physiol.* 189(3):275-284 (2001).  
Lopez-Coviella, I., et al., *Science* 289(5477):313-316 (2000).  
Miller, A.F., et al., *J. Biol. Chem.* 275(24):17937-17945 (2000).

#### BMP9 (GDF2) Antibody (N-term) - Citations

- [Reduced circulating BMP10 and BMP9 and elevated endoglin are associated with disease severity, decompensation and pulmonary vascular syndromes in patients with cirrhosis.](#)
- [Autocrine bone morphogenetic protein-9 signals through activin receptor-like kinase-2/Smad1/Smad4 to promote ovarian cancer cell proliferation.](#)

