

TGFB1 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21052a

Specification

TGFB1 Antibody - Product Information

Application	WB,E
Primary Accession	P01137
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	44341

TGFB1 Antibody - Additional Information

Gene ID 7040

Other Names

Transforming growth factor beta-1,
TGF-beta-1, Latency-associated peptide,
LAP, TGFB1, TGFB

Target/Specificity

This TGFB1 antibody is generated from a rabbit immunized with a recombinant protein of human TGFB1.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


Precautions

TGFB1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TGFB1 Antibody - Protein Information



Western blot analysis of lysates from K562, L929 cell line and rat lung tissue(from left to right), using TGFB1 Antibody(Cat. #AP21052a). AP21052a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.



Anti-TGFB1 Antibodyat 1:2000 dilution + A549 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit

Name TGFB1 ([HGNC:11766](#))

Synonyms TGFB

Function

Transforming growth factor beta-1 proprotein: Precursor of the Latency-associated peptide (LAP) and Transforming growth factor beta-1 (TGF-beta-1) chains, which constitute the regulatory and active subunit of TGF-beta-1, respectively.

Cellular Location

[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix

Tissue Location

Highly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158)

IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 44 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

TGFB1 Antibody - Background

Multifunctional protein that controls proliferation, differentiation and other functions in many cell types. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other growth factors. It plays an important role in bone remodeling as it is a potent stimulator of osteoblastic bone formation, causing chemotaxis, proliferation and differentiation in committed osteoblasts.

TGFB1 Antibody - References

Derynck R.,et al.Nucleic Acids Res. 15:3188-3189(1987).
Derynck R.,et al.Nature 316:701-705(1985).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

TGFB1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)