

(Mouse) Sox2 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21154a

Specification

(Mouse) Sox2 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P48432
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Calculated MW	34454

(Mouse) Sox2 Antibody (N-term) - Additional Information

Gene ID 20674

Other Names

Transcription factor SOX-2, Sox2, Sox-2

Target/Specificity

This Mouse Sox2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 34-68 amino acids from the N-terminal region of mouse Sox2.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

(Mouse) Sox2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.



Anti-(Mouse) Sox2 Antibody (N-term) at 1:1000 dilution + NCCIT whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size :34 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

(Mouse) Sox2 Antibody (N-term) - Background

Transcription factor that forms a trimeric complex with OCT4 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206. Critical for early embryogenesis and for embryonic stem cell pluripotency. May function as a switch in neuronal development. Downstream SRRT target that mediates the promotion of neural stem cell self-renewal. Keeps neural cells undifferentiated by counteracting the activity of proneural proteins and suppresses neuronal differentiation (By similarity).

(Mouse) Sox2 Antibody (N-term) - References

(Mouse) Sox2 Antibody (N-term) - Protein Information**Name** Sox2**Synonyms** Sox-2**Function**

Transcription factor that forms a trimeric complex with POU5F1 (OCT3/4) on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206 (PubMed:15863505, PubMed:17097055, PubMed:19740739). Binds to the proximal enhancer region of NANOG (PubMed:15863505). Critical for early embryogenesis and for embryonic stem cell pluripotency (By similarity). Downstream SRRT target that mediates the promotion of neural stem cell self-renewal (PubMed:22198669). Keeps neural cells undifferentiated by counteracting the activity of proneural proteins and suppresses neuronal differentiation (By similarity). May function as a switch in neuronal development (By similarity).

Cellular Location

Nucleus

{ECO:0000255|PROSITE-ProRule:PRU00267, ECO:0000269|PubMed:17097055, ECO:0000269|PubMed:32127020}

Tissue Location

Expressed in the cochlea (at protein level) (PubMed:32127020). Expressed in the brain and retina (PubMed:7590241, PubMed:15863505). A very low level of expression is seen in the stomach and lung (PubMed:7590241, PubMed:15863505). Expressed in the kidney (PubMed:15863505).

Yuan H.,et al.Genes Dev. 9:2635-2645(1995).

Yuan H.,et al.Submitted (AUG-1998) to the EMBL/GenBank/DDBJ databases.

Collignon J.,et al.Development 122:509-520(1996).

Tsuruzoe S.,et al.Biochem. Biophys. Res. Commun. 351:920-926(2006).

Takahashi K.,et al.Cell 126:663-676(2006).

**(Mouse) Sox2 Antibody (N-term) -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)