

NANOG Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21336c

Specification

NANOG Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q9H9S0
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Calculated MW	34620

NANOG Antibody (Center) - Additional Information

Gene ID 79923

Other Names

Homeobox protein NANOG, Homeobox transcription factor Nanog, hNanog, NANOG

Target/Specificity

This NANOG antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 155-189 amino acids from the Central region of human NANOG.

Dilution

WB~~1:2000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NANOG Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.



Anti-NANOG Antibody (Center) at 1:2000 dilution + human brain lysates
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution
Predicted band size : 35 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-NANOG Antibody (Center) at 1:2000 dilution + human brain lysates
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L),

NANOG Antibody (Center) - Protein Information**Name** NANOG**Function**

Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. Acts as a transcriptional activator or repressor. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3'. Binds to the POU5F1/OCT4 promoter (PubMed:25825768). Able to autorepress its expression in differentiating (ES) cells: binds to its own promoter following interaction with ZNF281/ZFP281, leading to recruitment of the NuRD complex and subsequent repression of expression. When overexpressed, promotes cells to enter into S phase and proliferation.

Cellular Location

Nucleus
{ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000269|PubMed:15983365}

Tissue Location

Expressed in testicular carcinoma and derived germ cell tumors (at protein level). Expressed in fetal gonads, ovary and testis. Also expressed in ovary teratocarcinoma cell line and testicular embryonic carcinoma. Not expressed in many somatic organs and oocytes.

NANOG Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

Peroxidase conjugated at 1/10000 dilution
Predicted band size : 35 kDa
Blocking/Dilution buffer: 5% NFDN/TBST.

NANOG Antibody (Center) - Background

Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. Acts as a transcriptional activator or repressor. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3'. Able to autorepress its expression in differentiating (ES) cells: binds to its own promoter following interaction with ZNF281/ZFP281, leading to recruitment of the NuRD complex and subsequent repression of expression. When overexpressed, promotes cells to enter into S phase and proliferation.

NANOG Antibody (Center) - References

Mitsui K.,et al.Cell 113:631-642(2003).
Clark A.T.,et al.Stem Cells 22:169-179(2004).
Kim J.S.,et al.Exp. Mol. Med. 37:601-607(2005).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)