

RAC2 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21363b

Specification

RAC2 Antibody (C-term) - Product Information

Application	IF, WB,E
Primary Accession	P15153
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG

RAC2 Antibody (C-term) - Additional Information

Gene ID 5880

Other Names

Ras-related C3 botulinum toxin substrate 2, GX, Small G protein, p21-Rac2, RAC2

Target/Specificity

This RAC2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 178-192 amino acids from the C-terminal region of human RAC2.

Dilution

IF~~1:25

WB~~1:2000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RAC2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized NIH/3T3 (mouse embryonic fibroblast cell line) cells labeling RAC2 with AP21363b at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG (NK179883) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm staining on NIH/3T3 cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red).The nuclear counter stain is DAPI (blue).

RAC2 Antibody (C-term) - Protein Information

Name RAC2

Function


Plasma membrane-associated small GTPase which cycles between an active GTP-bound and inactive GDP-bound state. In active state binds to a variety of effector proteins to regulate cellular responses, such as secretory processes, phagocytosis of apoptotic cells and epithelial cell polarization. Augments the production of reactive oxygen species (ROS) by NADPH oxidase.

Cellular Location

Cytoplasm. Note=Membrane-associated when activated

Tissue Location

Hematopoietic specific.



All lanes : Anti-RAC2 Antibody (C-term) at 1:2000 dilution
Lane 1: 293 whole cell lysate
Lane 2: Caco2 whole cell lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 21 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

RAC2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RAC2 Antibody (C-term) - Background

Plasma membrane-associated small GTPase which cycles between an active GTP-bound and inactive GDP-bound state. In active state binds to a variety of effector proteins to regulate cellular responses, such as secretory processes, phagocytosis of apoptotic cells and epithelial cell polarization. Augments the production of reactive oxygen species (ROS) by NADPH oxidase.

RAC2 Antibody (C-term) - References

Didsbury J.,et al.J. Biol. Chem. 264:16378-16382(1989).
Puhl H.L. III,et al.Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.
Collins J.E.,et al.Genome Biol. 5:R84.1-R84.11(2004).
Kalline N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Dunham I.,et al.Nature 402:489-495(1999).