

CASP14 Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21517a

Specification

CASP14 Antibody (N-Term) - Product Information

Application	WB,E
Primary Accession	P31944
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Calculated MW	27680

CASP14 Antibody (N-Term) - Additional Information

Gene ID 23581

Other Names

Caspase-14, CASP-14, 3422-, Caspase-14 subunit p19, Caspase-14 subunit p10, CASP14

Target/Specificity

This CASP14 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 59-93 amino acids from human CASP14.

Dilution

WB~~1:2000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CASP14 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.



All lanes : Anti-CASP14 Antibody (N-Term) at 1:2000 dilution Lane 1: A431 whole cell lysates Lane 2: MCF-7 whole cell lysates Lane 3: Jurkat whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 29 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

CASP14 Antibody (N-Term) - Background

Believed to be a non-apoptotic caspase which is involved in epidermal differentiation. Seems to play a role in keratinocyte differentiation and cornification. Probably regulates maturation of the epidermis by proteolytically processing filaggrin (By similarity).

CASP14 Antibody (N-Term) - References

Eckhart L.,et al.Biochem. Biophys. Res. Commun. 277:655-659(2000).
Pistritto G.,et al.Cell Death Differ. 9:995-1006(2002).
Rasmussen H.H.,et al.Electrophoresis 13:960-969(1992).
Lippens S.,et al.Cell Death Differ.

CASP14 Antibody (N-Term) - Protein Information

7:1218-1224(2000).

Chien A.J., et al. Biochem. Biophys. Res. Commun. 296:911-917(2002).

Name CASP14**Function**

Non-apoptotic caspase involved in epidermal differentiation. Is the predominant caspase in epidermal stratum corneum (PubMed:15556625). Seems to play a role in keratinocyte differentiation and is required for cornification. Regulates maturation of the epidermis by proteolytically processing filaggrin (By similarity). In vitro has a preference for the substrate [WY]-X-X-D motif and is active on the synthetic caspase substrate WEHD-ACF (PubMed:16854378, PubMed:19960512). Involved in processing of prosaposin in the epidermis (By similarity). May be involved in retinal pigment epithelium cell barrier function (PubMed:25121097). Involved in DNA degradation in differentiated keratinocytes probably by cleaving DFFA/ICAD leading to liberation of DFFB/CAD (PubMed:24743736).

Cellular Location

Cytoplasm. Nucleus

Tissue Location

Expressed in keratinocytes of adult skin suprabasal layers (from spinous layers to the stratum granulosum and stratum corneum) (at protein level). Expressed in keratinocytes of hair shaft and sebaceous glands (at protein level). In psoriatic skin only expressed at very low levels (PubMed:11175259). The p17/10 mature form is expressed in epidermis stratum corneum, the p20/p8 intermediate form in epidermis upper granular cells of the stratum granulosum (PubMed:22825846).

CASP14 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)