

NCSTN Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21573a

Specification

NCSTN Antibody (N-Term) - Product Information

Application	WB,E
Primary Accession	Q92542
Reactivity	Human, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Calculated MW	78411

NCSTN Antibody (N-Term) - Additional Information

Gene ID 23385

Other Names

Nicastrin, NCSTN, KIAA0253

Target/Specificity

This NCSTN antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 99-133 amino acids from human NCSTN.

Dilution

WB~~1:8000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


Precautions

NCSTN Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

NCSTN Antibody (N-Term) - Protein Information



All lanes : Anti-NCSTN Antibody (N-Term) at 1:2000 dilution Lane 1: Jurkat whole cell lysates Lane 2: A431 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 78 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-NCSTN Antibody (N-Term) at 1:8000 dilution Lane 1: HL-60 whole cell lysates Lane 2: HeLa whole cell lysates Lane 3: C6 whole cell lysates

Name NCSTN

Synonyms KIAA0253

Function

Essential subunit of the gamma-secretase complex, an endoprotease complex that catalyzes the intramembrane cleavage of integral membrane proteins such as Notch receptors and APP (amyloid- beta precursor protein) (PubMed:10993067, PubMed:12679784, PubMed:25043039, PubMed:26280335, PubMed:30598546, PubMed:30630874). The gamma-secretase complex plays a role in Notch and Wnt signaling cascades and regulation of downstream processes via its role in processing key regulatory proteins, and by regulating cytosolic CTNNB1 levels.

Cellular Location

Membrane; Single-pass type I membrane protein. Cytoplasmic vesicle membrane; Single-pass type I membrane protein. Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

Tissue Location

Detected in brain (at protein level) (PubMed:10993067). Widely expressed (PubMed:11396676)

3: C6 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 78 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

NCSTN Antibody (N-Term) - Background

Essential subunit of the gamma-secretase complex, an endoprotease complex that catalyzes the intramembrane cleavage of integral membrane proteins such as Notch receptors and APP (beta- amyloid precursor protein). It probably represents a stabilizing cofactor required for the assembly of the gamma-secretase complex.

NCSTN Antibody (N-Term) - References

Yu G.,et al.Nature 407:48-54(2000).
Clark H.F.,et al.Genome Res. 13:2265-2270(2003).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

NCSTN Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)