

RAP2A Antibody (C-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21722b

Specification

RAP2A Antibody (C-Term) - Product Information

Application	WB,E
Primary Accession	P10114
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Calculated MW	20615

RAP2A Antibody (C-Term) - Additional Information

Gene ID 5911

Other Names

Ras-related protein Rap-2a, RbBP-30, RAP2A

Target/Specificity

This RAP2A antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 124-159 amino acids from human RAP2A.

Dilution

WB~~1:2000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RAP2A Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.



Anti-RAP2A Antibody (C-Term) at 1:2000 dilution + A431 whole cell lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 21 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

RAP2A Antibody (C-Term) - Background

Small GTP-binding protein which cycles between a GDP- bound inactive and a GTP-bound active form. In its active form interacts with and regulates several effectors including MAP4K4, MINK1 and TNIK. Part of a signaling complex composed of NEDD4, RAP2A and TNIK which regulates neuronal dendrite extension and arborization during development. More generally, it is part of several signaling cascades and may regulate cytoskeletal rearrangements, cell migration, cell adhesion and cell spreading.

RAP2A Antibody (C-Term) - References

Pizon V.,et al.Oncogene 3:201-204(1988).
Fan Z.S.,et al.Submitted (NOV-1999) to the EMBL/GenBank/DDBJ databases.

RAP2A Antibody (C-Term) - Protein Information**Name** RAP2A**Function**

Small GTP-binding protein which cycles between a GDP-bound inactive and a GTP-bound active form. In its active form interacts with and regulates several effectors including MAP4K4, MINK1 and TNIK. Part of a signaling complex composed of NEDD4, RAP2A and TNIK which regulates neuronal dendrite extension and arborization during development. More generally, it is part of several signaling cascades and may regulate cytoskeletal rearrangements, cell migration, cell adhesion and cell spreading.

Cellular Location

Recycling endosome membrane; Lipid-anchor; Cytoplasmic side. Midbody. Note=May also localize to the Golgi (PubMed:7962206) and the gelatinase-containing granules of neutrophils (PubMed:8391995). Colocalizes with RASGEF1B to midbody at telophase (PubMed:23894443).

Puhl H.L. III,et al.Submitted (MAR-2002) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Dunham A.,et al.Nature 428:522-528(2004).

RAP2A Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)