

SQSTM1 (p62) Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2183B

Specification

SQSTM1 (p62) Antibody (C-term) - Product Information

Application	IF, WB, IHC-P,E
Primary Accession	Q13501
Other Accession	Q08623 , Q64337
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	317-346

SQSTM1 (p62) Antibody (C-term) - Additional Information

Gene ID 8878

Other Names

Sequestosome-1, EBI3-associated protein of 60 kDa, EBIAP, p60, Phosphotyrosine-independent ligand for the Lck SH2 domain of 62 kDa, Ubiquitin-binding protein p62, SQSTM1, ORCA, OSIL

Target/Specificity

This SQSTM1 (p62) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 317-346 amino acids of human SQSTM1 (p62).

Dilution


IF~~1:50~100
WB~~1:2000
IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2



Fluorescent image of U251 cells stained with SQSTM1 (p62) (C-term) antibody. U251 cells were treated with Chloroquine (50 μ M, 16h), then fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.2%, 30 min). Cells were then incubated with AP2183b SQSTM1 (p62) (C-term) primary antibody (1:200, 2 h at room temperature). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:1000, 1h). Nuclei were counterstained with Hoechst 33342 (blue) (10 μ g/ml, 5 min). SQSTM1 (p62) immunoreactivity is localized to autophagic vacuoles in the cytoplasm of U251 cells, supported by Human Protein Atlas Data (<http://www.proteinatlas.org/ENSG00000161011>).

weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SQSTM1 (p62) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.



SQSTM1 (p62) Antibody (C-term) - Protein Information

Name SQSTM1


Synonyms ORCA, OSIL

Function

Autophagy receptor required for selective macroautophagy (aggrephagy). Functions as a bridge between polyubiquitinated cargo and autophagosomes. Interacts directly with both the cargo to become degraded and an autophagy modifier of the MAP1 LC3 family (PubMed:16286508, PubMed:20168092, PubMed:24128730, PubMed:28404643, PubMed:22622177). Along with WDFY3, involved in the formation and autophagic degradation of cytoplasmic ubiquitin-containing inclusions (p62 bodies, ALIS/aggresome-like induced structures). Along with WDFY3, required to recruit ubiquitinated proteins to PML bodies in the nucleus (PubMed:24128730, PubMed:20168092). May regulate the activation of NFKB1 by TNF-alpha, nerve growth factor (NGF) and interleukin-1. May play a role in titin/TTN downstream signaling in muscle cells. May regulate signaling cascades through ubiquitination. Adapter that mediates the



Immunofluorescence staining of Autophagy SQSTM1 (p62) Antibody (C-term) (Cat# AP2183b) on Methanol-fixed and PFA fixed HeLa cells. Data courtesy of Dr. Eeva-Liisa Eskelinen, University of Helsinki, Finland.




All lanes : Anti-SQSTM1 (p62) Antibody (C-term) at 1:2000 dilution Lane 1: A549 whole cell lysate Lane 2: C2C12 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 48 kDa Blocking/Dilution buffer: 5% NFDM/TBST.


interaction between TRAF6 and CYLD (By similarity). May be involved in cell differentiation, apoptosis, immune response and regulation of K(+) channels. Involved in endosome organization by retaining vesicles in the perinuclear cloud: following ubiquitination by RNF26, attracts specific vesicle-associated adapters, forming a molecular bridge that restrains cognate vesicles in the perinuclear region and organizes the endosomal pathway for efficient cargo transport (PubMed:27368102). Promotes relocalization of 'Lys-63'-linked ubiquitinated STING1 to autophagosomes (PubMed:29496741). Acts as an activator of the NFE2L2/NRF2 pathway via interaction with KEAP1: interaction inactivates the BCR(KEAP1) complex, promoting nuclear accumulation of NFE2L2/NRF2 and subsequent expression of cytoprotective genes (PubMed:20452972, PubMed:28380357).

Cellular Location

Cytoplasm, cytosol. Late endosome. Lysosome. Cytoplasmic vesicle, autophagosome. Nucleus. Endoplasmic reticulum. Nucleus, PML body. Cytoplasm, myofibril, sarcomere. Note=In cardiac muscle, localizes to the sarcomeric band (By similarity). Commonly found in inclusion bodies containing polyubiquitinated protein aggregates. In neurodegenerative diseases, detected in Lewy bodies in Parkinson disease, neurofibrillary tangles in Alzheimer disease, and HTT aggregates in Huntington disease. In protein aggregate diseases of the liver, found in large amounts in Mallory bodies of alcoholic and nonalcoholic steatohepatitis, hyaline bodies in hepatocellular carcinoma, and in SERPINA1 aggregates Enriched in Rosenthal fibers of pilocytic astrocytoma. In the cytoplasm, observed in both membrane-free ubiquitin-containing protein aggregates (sequestosomes) and membrane-surrounded autophagosomes Colocalizes with TRIM13 in the perinuclear endoplasmic reticulum. Co- localizes with



All lanes : Anti-SQSTM1 (p62) Antibody (C-term) at 1:2000 dilution Lane 1: Jurkat whole cell lysate Lane 2: THP-1 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 48 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AP2183b staining SQSTM1 in Human prostate tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

TRIM5 in cytoplasmic bodies. When nuclear export is blocked by treatment with leptomycin B, accumulates in PML bodies


Tissue Location

Ubiquitously expressed.

SQSTM1 (p62) Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

SQSTM1 (p62) Antibody (C-term) - Background

SQSTM1/p62 is an adapter protein which binds ubiquitin and may regulate the activation of NF κ B1 by TNF-alpha, nerve growth factor (NGF) and interleukin-1. This protein may play a role in titin/TTN downstream signaling in muscle cells, and may also regulate signaling cascades through ubiquitination. This protein is involved in cell differentiation, apoptosis, immune response and regulation of K(+) channels. SQSTM1/p62 also appears to play a role in macroautophagic removal of intracellular protein aggregates. Cellular depletion studies of SQSTM1/p62 have indicated a role for association with LC3 and aggregate proteins in order to facilitate normal formation of the autophagosome.

SQSTM1 (p62) Antibody (C-term) - References

References for protein:

1. Seibenhener, M.L., et al., Mol. Cell. Biol. 24(18):8055-8068 (2004).
2. Eekhoff, E.W., et al., Arthritis Rheum. 50(5):1650-1654 (2004).
3. Brajenovic, M., et al., J. Biol. Chem.

279(13):12804-12811 (2004).

4. Kuusisto, E., et al., J. Neuropathol. Exp. Neurol. 62(12):1241-1253 (2003).

5. Johnson-Pais, T.L., et al., J. Bone Miner. Res. 18(10):1748-1753 (2003).

References for U251 cell line:

1. Westermarck B.; Pontén J.; Hugosson R. (1973). "Determinants for the establishment of permanent tissue culture lines from human gliomas". Acta Pathol Microbiol Scand A. 81:791-805. [PMID: 4359449].

2. Pontén, J., Westermarck B. (1978). "Properties of Human Malignant Glioma Cells in Vitro". Medical Biology 56: 184-193. [PMID: 359950].

3. Geng Y.; Kohli L.; Klocke B.J.; Roth K.A. (2010). "Chloroquine-induced autophagic vacuole accumulation and cell death in glioma cells is p53 independent". Neuro Oncol. 12(5): 473-481. [PMID: 20406898].

SQSTM1 (p62) Antibody (C-term) - Citations

- [Rapamycin induces megakaryocytic differentiation through increasing autophagy in Dami cells](#)
- [Enteritidis Effector AvrA Suppresses Autophagy by Reducing Beclin-1 Protein](#)
- [Axonal autophagosome maturation defect through failure of ATG9A sorting underpins pathology in AP-4 deficiency syndrome](#)
- [The p53 inactivators pifithrin-μ and pifithrin-α mitigate TBI-induced neuronal damage through regulation of oxidative stress, neuroinflammation, autophagy and mitophagy](#)
- [Helicobacter pylori cholesterol glucosylation modulates autophagy for increasing intracellular survival in macrophages](#)
- [Genistein and Myd88 Activate Autophagy in High Glucose-Induced Renal Podocytes In Vitro](#)
- [Honokiol inhibits in vitro and in vivo growth of oral squamous cell carcinoma through induction of apoptosis, cell cycle arrest and autophagy](#)
- [Up-regulation of autophagy is a mechanism of resistance to chemotherapy and can be inhibited by pantoprazole to increase drug sensitivity](#)
- [Deletion of the BH3-only protein Noxa alters electrographic seizures but does not protect against hippocampal damage after status epilepticus in mice](#)
- [Role of Autophagy as a Survival Mechanism for Hypoxic Cells in Tumors](#)
- [Interference with HMGB1 increases the sensitivity to chemotherapy drugs by inhibiting HMGB1-mediated cell autophagy and inducing cell apoptosis](#)
- [Effect of pantoprazole to enhance activity of docetaxel against human tumour xenografts by inhibiting autophagy](#)
- [Intestinal epithelial vitamin D receptor deletion leads to defective autophagy in colitis](#)
- [Inhibition of Intracellular Clusterin Attenuates Cell Death in Nephropathic Cystinosis](#)
- [A novel sulindac derivative inhibits lung adenocarcinoma cell growth through suppression of Akt/mTOR signaling and induction of autophagy](#)
- [Potent obatoclax cytotoxicity and activation of triple death mode killing across infant acute lymphoblastic leukemia](#)
- [p62/SQSTM1 prominently accumulates in renal proximal tubules in nephropathic cystinosis](#)
- [Curcumin induces autophagy to protect vascular endothelial cell survival from oxidative stress damage](#)
- [Increased hippocampal accumulation of autophagosomes predicts short-term recognition memory impairment in aged mice](#)

- [Induction of an incomplete autophagic response by cancer-preventive geranylgeranoic acid \(GGA\) in a human hepatoma-derived cell line.](#)
- [Overexpression of the autophagic beclin-1 protein clears mutant ataxin-3 and alleviates Machado-Joseph disease.](#)
- [Autophagy negatively regulates keratinocyte inflammatory responses via scaffolding protein p62/SQSTM1.](#)
- [Roles of SIRT1 in the acute and restorative phases following induction of inflammation.](#)
- [Invasion and multiplication of Helicobacter pylori in gastric epithelial cells and implications for antibiotic resistance.](#)
- [Epidermal growth factor reduces autophagy in intestinal epithelium and in the rat model of necrotizing enterocolitis.](#)
- [Autophagy induction with RAD001 enhances chemosensitivity and radiosensitivity through Met inhibition in papillary thyroid cancer.](#)
- [Helicobacter pylori impairs murine dendritic cell responses to infection.](#)
- [Transcription factor GATA4 inhibits doxorubicin-induced autophagy and cardiomyocyte death.](#)
- [Absence of autophagy results in reactive oxygen species-dependent amplification of RLR signaling.](#)
- [Impaired protein aggregate handling and clearance underlie the pathogenesis of p97/VCP-associated disease.](#)