

ARHGEF7 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP22153b

Specification

ARHGEF7 Antibody (C-Term) - Product Information

Application WB,E
Primary Accession Other Accession O9ES28

Reactivity Human, Mouse

Predicted Mouse
Host Rabbit
Clonality polyclonal
Isotype Rabbit Ig
Calculated MW 90012

ARHGEF7 Antibody (C-Term) - Additional Information

Gene ID 8874

Other Names

Rho guanine nucleotide exchange factor 7, Beta-Pix, COOL-1, PAK-interacting exchange factor beta, p85, ARHGEF7, COOL1, KIAA0142, P85SPR, PAK3BP, PIXB

Target/Specificity

This ARHGEF7 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 654-688 amino acids from human ARHGEF7.

Dilution

WB~~1:2000


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions




All lanes: Anti-ARHGEF7 Antibody (C-Term) at 1:2000 dilution Lane 1: Jurkat whole cell lysate Lane 2: MCF-7 whole cell lysate Lane 3: NIH/3T3 whole cell lysate Lane 4: ZR-75-1 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 90 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

ARHGEF7 Antibody (C-Term) - Background

Acts as a RAC1 guanine nucleotide exchange factor (GEF) and can induce membrane ruffling. Functions in cell migration, attachment and cell spreading. Promotes targeting of RAC1 to focal adhesions (By similarity). May function as a positive regulator of apoptosis. Downstream of NMDA receptors and CaMKK-CaMK1 signaling cascade, promotes the formation of spines and synapses in hippocampal neurons.

ARHGEF7 Antibody (C-Term) - References

Nagase T.,et al.DNA Res. 2:167-174(1995). Bagrodia S.,et al.J. Biol. Chem. 273:23633-23636(1998).





Tel: 858.875.1900 Fax: 858.622.0609

ARHGEF7 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

8:399-399(2007). Dunham A., et al. Nature 428:522-528(2004). Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBI databases.

Bechtel S., et al. BMC Genomics

ARHGEF7 Antibody (C-Term) - Protein Information

Name ARHGEF7

Synonyms COOL1, KIAA0142, P85SPR, PAK3BP, PIXB

Function

Acts as a RAC1 quanine nucleotide exchange factor (GEF) and can induce membrane ruffling. Functions in cell migration, attachment and cell spreading. Promotes targeting of RAC1 to focal adhesions (By similarity). May function as a positive regulator of apoptosis. Downstream of NMDA receptors and CaMKK-CaMK1 signaling cascade, promotes the formation of spines and synapses in hippocampal neurons.

Cellular Location

Cell junction, focal adhesion. Cell projection, ruffle. Cytoplasm, cell cortex Cell projection, lamellipodium. Note=Detected at cell adhesions. A small proportion is detected at focal adhesions

ARHGEF7 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cvtometv
- Cell Culture