

**ACSL3 (FACL3) Antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2535A**

**Specification**

**ACSL3 (FACL3) Antibody (N-term) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">095573</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	80420
Antigen Region	1-30

**ACSL3 (FACL3) Antibody (N-term) - Additional Information**

**Gene ID** 2181

**Other Names**

Long-chain-fatty-acid--CoA ligase 3,  
Long-chain acyl-CoA synthetase 3, LACS 3,  
ACSL3, ACS3, FACL3, LACS3

**Target/Specificity**

This ACSL3 (FACL3) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human ACSL3 (FACL3).

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100


**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.


**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**



Western blot analysis of anti-FACL3 Antibody (N-term) (Cat.#AP2535a) in 293 cell line lysates (35ug/lane). FACL3 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

ACSL3 (FACL3) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### **ACSL3 (FACL3) Antibody (N-term) - Protein Information**

**Name** ACSL3 ([HGNC:3570](#))

**Synonyms** ACS3, FACL3, LACS3

#### **Function**

Acyl-CoA synthetases (ACSL) activates long-chain fatty acids for both synthesis of cellular lipids, and degradation via beta-oxidation (PubMed:<a href="http://www.uniprot.org/citations/22633490" target="\_blank">22633490</a>). Required for the incorporation of fatty acids into phosphatidylcholine, the major phospholipid located on the surface of VLDL (very low density lipoproteins) (PubMed:<a href="http://www.uniprot.org/citations/18003621" target="\_blank">18003621</a>). Has mainly an anabolic role in energy metabolism. Mediates hepatic lipogenesis. Preferentially uses myristate, laurate, arachidonate and eicosapentaenoate as substrates. Both isoforms exhibit the same level of activity (By similarity).

#### **Cellular Location**

Mitochondrion outer membrane; Single-pass type III membrane protein. Peroxisome membrane; Single-pass type III membrane protein. Microsome membrane; Single-pass type III membrane protein. Endoplasmic reticulum membrane; Single-pass type III membrane protein

#### **ACSL3 (FACL3) Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **ACSL3 (FACL3) Antibody (N-term) - Background**

An initial reaction in fatty acid metabolism in eukaryotic cells is activation of fatty acids catalyzed by acyl-CoA synthetase. FACL3 (fatty acid CoA ligase, long-chain 3) is identified as member of the acyl-CoA synthetase (ACS) family by PCR of rat brain cDNAs using primers based on the conserved region of the ACS protein. The 720-amino acid rat protein preferentially utilizes myristate, laurate, arachidonate, and eicosapentaenoate, and is expressed primarily in brain. The predicted 720-amino acid FACL3 human protein is 92% identical to that of rat.

#### **ACSL3 (FACL3) Antibody (N-term) - References**

Genomics 42:180-181(1997).  
Gene 278:185-192(2001).