

### **SULT1C1** Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2602b

## **Specification**

#### **SULT1C1** Antibody (Center) - Product Information

Application WB,E Primary Accession 000338 NP 789795 Other Accession Human, Mouse Reactivity Host Rabbit Clonality **Polyclonal** Isotype Rabbit Ig Calculated MW 34880 Antigen Region 95-127

SULT1C1 Antibody (Center) - Additional Information

#### **Gene ID 6819**

#### **Other Names**

Sulfotransferase 1C2, ST1C2, 282-, Sulfotransferase 1C1, SULT1C#1, humSULTC2, SULT1C2, SULT1C1

#### Target/Specificity

This SULT1C1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 95-127 amino acids from the Central region of human SULT1C1.

## **Dilution**

WB~~1:1000

# **Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

# **Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

SULT1C1 Antibody (Center) is for research



The anti-SULT1C1 Pab (Cat. #AP2602b) is used in Western blot to detect SULT1C1 in mouse kidney tissue lysate.

# SULT1C1 Antibody (Center) - Background

The 296-amino acid human SULTC1 protein, so named on the basis of its significant homology to a rat hepatic cytosolic sulfotranserase ST1C1, catalyzes the sulfate conjugation of many drugs, xenobiotic compounds, hormones, and neurotransmitters, and may be involved in the activation of carcinogenic hydroxylamines. This enzyme also shows activity towards p-nitrophenol and N-hydroxy-2-acetylamino-fluorene (N-OH-2AAF). SULT1C1 is expressed as a 1.4-kb mRNA in adult human stomach, kidney, and thyroid, and in fetal kidney and liver. By functional characterization of recombinant protein, it has been shown that SULT1C1 catalyzes the sulfonation of p-nitrophenol and N-hydroxy-2-acetylaminofluorene, but not dopamine, 17-beta-estradiol, or dehydroepiandrosterone.



use only and not for use in diagnostic or therapeutic procedures.

**SULT1C1** Antibody (Center) - Protein Information

Name SULT1C2

**Synonyms** SULT1C1

#### **Function**

Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the sulfate conjugation of drugs, xenobiotic compounds, hormones, and neurotransmitters. May be involved in the activation of carcinogenic hydroxylamines. Shows activity towards p-nitrophenol and N-hydroxy-2-acetylamino-fluorene (N-OH-2AAF).

Cellular Location Cytoplasm.

## **Tissue Location**

Found in adult stomach, kidney and thyroid gland, and in fetal kidney and liver

# **SULT1C1** Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture