

**FZD1 Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2755C**

### Specification

#### FZD1 Antibody (Center) - Product Information

Application	WB, IHC-P, FC, E
Primary Accession	<a href="#">Q9UP38</a>
Other Accession	<a href="#">Q08463</a> , <a href="#">O70421</a>
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	71158
Antigen Region	367-396

#### FZD1 Antibody (Center) - Additional Information

**Gene ID** 8321

#### Other Names

Frizzled-1, Fz-1, hFz1, FzE1, FZD1

#### Target/Specificity

This FZD1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 367-396 amino acids from the Central region of human FZD1.

#### Dilution

WB~~1:1000  
IHC-P~~1:10~50  
FC~~1:10~50

#### Format



Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

FZD1 Antibody (Center) is for research use



Formalin-fixed and paraffin-embedded human prostate carcinoma tissue reacted with FZD1 antibody (Center) (Cat.#AP2755c), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

only and not for use in diagnostic or therapeutic procedures.

#### FZD1 Antibody (Center) - Protein Information

**Name** FZD1

#### Function


Receptor for Wnt proteins (PubMed:<a href = "http://www.uniprot.org/citations/10557084" target="\_blank">10557084</a>). Activated by WNT3A, WNT3, WNT1 and to a lesser extent WNT2, but apparently not by WNT4, WNT5A, WNT5B, WNT6, WNT7A or WNT7B (PubMed:<a href="http://www.uniprot.org/citations/10557084" target="\_blank">10557084</a>). Contradictory results showing activation by WNT7B have been described for mouse (By similarity). Functions in the canonical Wnt/beta-catenin signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/10557084" target="\_blank">10557084</a>). The canonical Wnt/beta-catenin signaling pathway leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes (PubMed:<a href="http://www.uniprot.org/citations/10557084" target="\_blank">10557084</a>). A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues (Probable).

#### Cellular Location

Cell membrane; Multi-pass membrane protein

#### Tissue Location

Expressed in adult heart, placenta, lung, kidney, pancreas, prostate, and ovary and in fetal lung and kidney



Flow cytometric analysis of NCI-H292 cells using FZD1 Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### FZD1 Antibody (Center) - Background

Members of the 'frizzled' family are 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The FZD1 protein contains a signal peptide, a cysteine-rich domain in the N-terminal extracellular region, 7 transmembrane domains, and a C-terminal PDZ domain-binding motif.

#### FZD1 Antibody (Center) - References

- Quelard,D., (er) PLoS ONE 3 (4), E1878 (2008)
- Hardie,W.D.,Am. J. Respir. Cell Mol. Biol. 37 (3), 309-321 (2007)
- Yang,L., J. Dermatol. Sci. 42 (2), 111-119 (2006)

## FZD1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)