

B2M Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2771b

Specification

B2M Antibody (C-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P61769
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	86-118

B2M Antibody (C-term) - Additional Information

Gene ID 567

Other Names

Beta-2-microglobulin, Beta-2-microglobulin form pl 53, B2M

Target/Specificity

This B2M antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 86-118 amino acids from the C-terminal region of human B2M.

Dilution

WB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

B2M Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.



B2M Antibody (C-term) (Cat. #AP2771b) western blot analysis in Jurkat cell line lysates (35ug/lane). This demonstrates the B2M antibody detected the B2M protein (arrow).



Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with B2M antibody (C-term) (Cat.#AP2771b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

B2M Antibody (C-term) - Protein Information

Name B2M ([HGNC:914](#))

Function

Component of the class I major histocompatibility complex (MHC). Involved in the presentation of peptide antigens to the immune system. Exogenously applied *M. tuberculosis* EsxA or EsxA-EsxB (or EsxA expressed in host) binds B2M and decreases its export to the cell surface (total protein levels do not change), probably leading to defects in class I antigen presentation (PubMed:25356553).

Cellular Location

Secreted. Cell surface. Note=Detected in serum and urine (PubMed:1336137, PubMed:7554280).
{ECO:0000269|PubMed:7554280, ECO:0000269|Ref.6}


B2M Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

B2M Antibody (C-term) - Citations

- [Overexpression of B2M and loss of ALK7 expression are associated with invasion, metastasis, and poor-prognosis of the pancreatic ductal adenocarcinoma.](#)



B2M Antibody (C-term) (Cat. #AP2771b) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

B2M Antibody (C-term) - Background

Beta-2-microglobulin is a serum protein found in association with the major histocompatibility complex (MHC) class I heavy chain on the surface of nearly all nucleated cells.

B2M Antibody (C-term) - References

Sakata,M., J. Mol. Biol. 382 (5), 1242-1255 (2008)
Huang,W.C., Clin. Cancer Res. 14 (17), 5341-5347 (2008)
Gattoni-Celli,S., Cancer Res. 52 (5), 1201-1204 (1992)