

EXO1 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2871a

Specification

EXO1 Antibody (N-term) - Product Information

Application	IF, WB,E
Primary Accession	Q9UQ84
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	1-30

EXO1 Antibody (N-term) - Additional Information

Gene ID 9156

Other Names

Exonuclease 1, hExo1, 31--, Exonuclease I, hExo1, EXO1, EXO1, HEX1

Target/Specificity

This EXO1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human EXO1.

Dilution

IF~~1:10~50

WB~~1:2000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

EXO1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.



Confocal immunofluorescent analysis of EXO1 Antibody (N-term) (Cat. #AP2871a) with A549 cell followed by Alexa Fluor® 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



All lanes : Anti-EXO1 Antibody (N-term) at 1:2000 dilution Lane 1: Jurkat whole cell lysate Lane 2: U-2OS whole cell lysate Lane 3: A549 whole cell lysate Lane 4: K562 whole cell lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),

EXO1 Antibody (N-term) - Protein Information**Name** EXO1**Synonyms** EXO1, HEX1**Function**

5'->3' double-stranded DNA exonuclease which may also possess a cryptic 3'->5' double-stranded DNA exonuclease activity. Functions in DNA mismatch repair (MMR) to excise mismatch-containing DNA tracts directed by strand breaks located either 5' or 3' to the mismatch. Also exhibits endonuclease activity against 5'-overhanging flap structures similar to those generated by displacement synthesis when DNA polymerase encounters the 5'-end of a downstream Okazaki fragment. Required for somatic hypermutation (SHM) and class switch recombination (CSR) of immunoglobulin genes. Essential for male and female meiosis.

Cellular Location

Nucleus Note=Colocalizes with PCNA to discrete nuclear foci in S-phase

Tissue Location

Highly expressed in bone marrow, testis and thymus. Expressed at lower levels in colon, lymph nodes, ovary, placenta, prostate, small intestine, spleen and stomach

Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 94 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

EXO1 Antibody (N-term) - Background

EXO1 is a protein with 5' to 3' exonuclease activity as well as an RNase H activity. This protein is similar to the *Saccharomyces cerevisiae* protein Exo1 which interacts with Msh2 and which is involved in mismatch repair and recombination.

EXO1 Antibody (N-term) - References

Tishkoff D.X., Amin N.S., Viars C.S. *Cancer Res.* 58:5027-5031(1998)
Lee B.-I., Nguyen L.H., *Nucleic Acids Res.* 30:942-949(2002)
Alam N.A., Gorman P., Jaeger E.E.M. *Cancer Genet. Cytogenet.* 147:121-127(2003)
Knudsen N.O., Nielsen F.C., Vinther L. *Nucleic Acids Res.* 35:2609-2619(2007)

EXO1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)