

HSPA8 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP2872b

Specification

HSPA8 Antibody (C-term) - Product Information

Application	IF, WB, IHC-P, FC, E
Primary Accession	P11142
Other Accession	P63018 , P63017 , P19378 , P19120 , A2Q0Z1
Reactivity Predicted	Human, Mouse, Bovine, Hamster, Horse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	539-569

HSPA8 Antibody (C-term) - Additional Information

Gene ID 3312

Other Names

Heat shock cognate 71 kDa protein, Heat shock 70 kDa protein 8, Lipopolysaccharide-associated protein 1, LAP-1, LPS-associated protein 1, HSPA8, HSC70, HSP73, HSPA10

Target/Specificity


This HSPA8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 539-569 amino acids from the C-terminal region of human HSPA8.

Dilution


IF~~1:10~50
WB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.



Confocal immunofluorescent analysis of HSPA8 Antibody (C-term)(Cat#AP2872b) with HeLa cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red).



Western blot analysis of lysates from rat H-4-II-E, mouse NIH/3T3 cell line (from left to right), using HSPA8 Antibody (C-term) (Cat. # AP2872b). AP2872b was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HSPA8 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.


HSPA8 Antibody (C-term) - Protein Information

Name HSPA8 ([HGNC:5241](#))


Function

Molecular chaperone implicated in a wide variety of cellular processes, including protection of the proteome from stress, folding and transport of newly synthesized polypeptides, activation of proteolysis of misfolded proteins and the formation and dissociation of protein complexes. Plays a pivotal role in the protein quality control system, ensuring the correct folding of proteins, the re-folding of misfolded proteins and controlling the targeting of proteins for subsequent degradation (PubMed:21150129, PubMed:21148293, PubMed:24732912, PubMed:27916661, PubMed:23018488). This is achieved through cycles of ATP binding, ATP hydrolysis and ADP release, mediated by co-chaperones (PubMed:21150129, PubMed:21148293, PubMed:24732912, PubMed:27916661).

1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



HSPA8 Antibody (C-term) (Cat. #AP2872b) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of HSPA8 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



HSPA8 Antibody (C-term) (Cat. #AP2872b) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

HSPA8 Antibody (C-term) - Background

HSPA8 belongs to the heat shock protein 70

target="_blank">27916661,
PubMed:<a href="http://www.uniprot.org/citations/23018488"
target="_blank">23018488,
PubMed:<a href="http://www.uniprot.org/citations/12526792"
target="_blank">12526792). The
co-chaperones have been shown to not only
regulate different steps of the ATPase cycle
of HSP70, but they also have an individual
specificity such that one co-chaperone may
promote folding of a substrate while
another may promote degradation
(PubMed:<a href="http://www.uniprot.org/citations/21150129"
target="_blank">21150129,
PubMed:<a href="http://www.uniprot.org/citations/21148293"
target="_blank">21148293,
PubMed:<a href="http://www.uniprot.org/citations/24732912"
target="_blank">24732912,
PubMed:<a href="http://www.uniprot.org/citations/27916661"
target="_blank">27916661,
PubMed:<a href="http://www.uniprot.org/citations/23018488"
target="_blank">23018488,
PubMed:<a href="http://www.uniprot.org/citations/12526792"
target="_blank">12526792). The
affinity of HSP70 for polypeptides is
regulated by its nucleotide bound state. In
the ATP-bound form, it has a low affinity for
substrate proteins. However, upon
hydrolysis of the ATP to ADP, it undergoes a
conformational change that increases its
affinity for substrate proteins. HSP70 goes
through repeated cycles of ATP hydrolysis
and nucleotide exchange, which permits
cycles of substrate binding and release. The
HSP70-associated co-chaperones are of
three types: J- domain co-chaperones
HSP40s (stimulate ATPase hydrolysis by
HSP70), the nucleotide exchange factors
(NEF) such as BAG1/2/3 (facilitate
conversion of HSP70 from the ADP-bound to
the ATP-bound state thereby promoting
substrate release), and the TPR domain
chaperones such as HOPX and STUB1
(PubMed:<a href="http://www.uniprot.org/citations/24318877"
target="_blank">24318877,
PubMed:<a href="http://www.uniprot.org/citations/27474739"
target="_blank">27474739,
PubMed:<a href="http://www.uniprot.org/ci

family which contains both heat-inducible and
constitutively expressed members. The latter
are called heat-shock cognate proteins. HSPA8
is a heat-shock cognate protein. This protein
binds to nascent polypeptides to facilitate
correct folding. The protein also functions as
an ATPase in the disassembly of
clathrin-coated vesicles during transport of
membrane components through the cell.

HSPA8 Antibody (C-term) - References

Tsukahara F., Yoshioka T. Mol. Pharmacol.
58:1257-1263(2000)
Egerton M., Moritz R.L. Biochem. Biophys. Res.
Commun. 224:666-674(1996)

tations/24121476"
target="_blank">24121476,
PubMed:<a href="http://www.uniprot.org/ci
tations/26865365"
target="_blank">26865365). Plays a
critical role in mitochondrial import, delivers
preproteins to the mitochondrial import
receptor TOMM70 (PubMed:<a href="http://
www.uniprot.org/citations/12526792"
target="_blank">12526792). Acts as a
repressor of transcriptional activation.
Inhibits the transcriptional coactivator
activity of CITED1 on Smad- mediated
transcription. Component of the
PRP19-CDC5L complex that forms an
integral part of the spliceosome and is
required for activating pre- mRNA splicing.
May have a scaffolding role in the
spliceosome assembly as it contacts all
other components of the core complex.
Binds bacterial lipopolysaccharide (LPS) and
mediates LPS-induced inflammatory
response, including TNF secretion by
monocytes (PubMed:<a href="http://www.u
niprot.org/citations/10722728"
target="_blank">10722728,
PubMed:<a href="http://www.uniprot.org/ci
tations/11276205"
target="_blank">11276205).
Participates in the ER-associated
degradation (ERAD) quality control pathway
in conjunction with J domain-containing
co-chaperones and the E3 ligase STUB1
(PubMed:<a href="http://www.uniprot.org/c
itations/23990462"
target="_blank">23990462). Interacts
with VGF-derived peptide TLQP-21
(PubMed:<a href="http://www.uniprot.org/c
itations/28934328"
target="_blank">28934328).

Cellular Location

Cytoplasm. Melanosome. Nucleus,
nucleolus. Cell membrane. Note=Localized
in cytoplasmic mRNP granules containing
untranslated mRNAs. Translocates rapidly
from the cytoplasm to the nuclei, and
especially to the nucleoli, upon heat shock

Tissue Location

Ubiquitous..

HSPA8 Antibody (C-term) - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)