

GSTP1 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2932b

Specification

GSTP1 Antibody (C-term) - Product Information

Application	IF, WB, FC, E
Primary Accession	P09211
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	23356
Antigen Region	165-192

GSTP1 Antibody (C-term) - Additional Information

Gene ID 2950

Other Names

Glutathione S-transferase P, GST class-pi,
GSTP1-1, GSTP1, FAES3, GST3

Target/Specificity

This GSTP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 165-192 amino acids from the C-terminal region of human GSTP1.

Dilution

IF~~1:10~50
WB~~1:1000
FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GSTP1 Antibody (C-term) is for research use only and not for use in diagnostic or



Confocal immunofluorescent analysis of GSTP1 Antibody (C-term)(Cat#AP2932b) with HeLa cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



GSTP1 Antibody (C-term) (Cat. #AP2932b) western blot analysis in K562, PC-3, Y79 cell line lysates (35ug/lane). This demonstrates the GSTP1 antibody detected the GSTP1 protein (arrow).

therapeutic procedures.

GSTP1 Antibody (C-term) - Protein Information

Name GSTP1 ([HGNC:4638](#))

Synonyms FAES3, GST3

Function

Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. Involved in the formation of glutathione conjugates of both prostaglandin A2 (PGA2) and prostaglandin J2 (PGJ2) (PubMed:9084911). Participates in the formation of novel hepxilin regioisomers (PubMed:21046276). Regulates negatively CDK5 activity via p25/p35 translocation to prevent neurodegeneration.


Cellular Location

Cytoplasm. Mitochondrion. Nucleus.
Note=The 83 N-terminal amino acids function as an uncleaved transit peptide, and arginine residues within it are crucial for mitochondrial localization

GSTP1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Flow cytometric analysis of wiDr cells using GSTP1 Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram) FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

GSTP1 Antibody (C-term) - Background

Glutathione S-transferases (GSTs) are a family of enzymes that play an important role in detoxification by catalyzing the conjugation of many hydrophobic and electrophilic compounds with reduced glutathione. Based on their biochemical, immunologic, and structural properties, the soluble GSTs are categorized into 4 main classes: alpha, mu, pi, and theta. This GST family member is a polymorphic gene encoding active, functionally different GSTP1 variant proteins that are thought to function in xenobiotic metabolism and play a role in susceptibility to cancer, and other diseases.

GSTP1 Antibody (C-term) - References

- Spurdle,A.B., et.al., Breast Cancer Res. Treat. (2009)
Agusa,T., et.al., Toxicol. Appl. Pharmacol. (2009)