

**Phospho-LC3C(S12) Antibody**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP3301a**

**Specification**

**Phospho-LC3C(S12) Antibody - Product Information**

|                      |                                                                                                      |
|----------------------|------------------------------------------------------------------------------------------------------|
| Application          | WB, DB, E                                                                                            |
| Primary Accession    | <a href="#">Q9H492</a>                                                                               |
| Other Accession      | <a href="#">Q6XVN8</a> , <a href="#">Q91VR7</a> ,<br><a href="#">Q2HJ23</a> , <a href="#">Q6NX90</a> |
| Reactivity Predicted | Human<br>Zebrafish, Bovine,<br>Mouse, Rat                                                            |
| Host                 | Rabbit                                                                                               |
| Clonality            | Polyclonal                                                                                           |
| Isotype              | Rabbit Ig                                                                                            |

**Phospho-LC3C(S12) Antibody - Additional Information**

**Gene ID** 84557

**Other Names**

Microtubule-associated proteins 1A/1B light chain 3A, Autophagy-related protein LC3 A, Autophagy-related ubiquitin-like modifier LC3 A, MAP1 light chain 3-like protein 1, MAP1A/MAP1B light chain 3 A, MAP1A/MAP1B LC3 A, Microtubule-associated protein 1 light chain 3 alpha, MAP1LC3A

**Target/Specificity**


This LC3C Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S12 of human LC3C.

**Dilution**


WB~~1:1000  
DB~~1:500

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.



Immunoblots of phosphorylated LC3 (phospho-LC3) in CHO cell culture. LC3 and LC3 S12A mutant vectors were transfected into CHO cells. The cell lysates were separated with SDS-PAGE and blotted with anti-phospho-LC3 S12 antibody. LC3 = microtubule-associated protein light chain-3; S12A = replacement of the amino acid position 12 serine of LC3 with alanine. WT = wildtype LC3-transfected cell lysates; S12A = LC3 S12A mutant-transfected cell lysates; Empty vector = vector with no LC3 gene. Molecular size: LC3-I = 16kDa, and LC3-II = 14 kDa



Immunoblots of SH-SY5Y cells treated with rapamycin for 1 h was probed with AP3301a. The data shows that treatment with rapamycin showed no significant change in level of LC3.



Immunoblots of SH-SY5Y cells treated with MPP+ for 24h was probed with AP3301a. The data shows that treatment with MPP+

### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### Precautions

Phospho-LC3C(S12) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### Phospho-LC3C(S12) Antibody - Protein Information

**Name** MAP1LC3A

### Function

Ubiquitin-like modifier involved in formation of autophagosomal vacuoles (autophagosomes) (PubMed:<a href="http://www.uniprot.org/citations/20713600" target="\_blank">20713600</a>, PubMed:<a href="http://www.uniprot.org/citations/24290141" target="\_blank">24290141</a>). While LC3s are involved in elongation of the phagophore membrane, the GABARAP/GATE-16 subfamily is essential for a later stage in autophagosome maturation (PubMed:<a href="http://www.uniprot.org/citations/20713600" target="\_blank">20713600</a>). Through its interaction with the reticulophagy receptor TEX264, participates in the remodeling of subdomains of the endoplasmic reticulum into autophagosomes upon nutrient stress, which then fuse with lysosomes for endoplasmic reticulum turnover (PubMed:<a href="http://www.uniprot.org/citations/31006538" target="\_blank">31006538</a>, PubMed:<a href="http://www.uniprot.org/citations/31006537" target="\_blank">31006537</a>).


### Cellular Location

Cytoplasm, cytoskeleton. Endomembrane system; Lipid-anchor. Cytoplasmic vesicle, autophagosome membrane; Lipid- anchor. Cytoplasmic vesicle, autophagosome. Note=LC3-II binds to the autophagic membranes


### Tissue Location

Most abundant in heart, brain, liver, skeletal

showed no significant change in level of LC3.



Something like SH-SY5Y cells expressing GFP-LC3-WT or-S12D treated with rapamycin or vehicle for 1h.



Dot blot analysis of Phospho-LC3 (APG8a) - S12 Antibody (Cat. #AP3301a) and Nonphospho-LC3 (APG8a) Antibody on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

### Phospho-LC3C(S12) Antibody - Background

MAP1A and MAP1B are microtubule-associated proteins which mediate the physical interactions between microtubules and components of the cytoskeleton. These proteins are involved in formation of autophagosomal vacuoles (autophagosomes). MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. MAP1LC3a is one of the light chain subunits and can associate with either MAP1A or MAP1B. The precursor molecule is cleaved by APG4B/ATG4B to form the cytosolic form,

muscle and testis but absent in thymus and peripheral blood leukocytes

### **Phospho-LC3C(S12) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form the membrane-bound form, LC3-II. Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole).

### **Phospho-LC3C(S12) Antibody - References**

Baehrecke EH. Nat Rev Mol Cell Biol. 6(6):505-10. (2005) Lum JJ, et al. Nat Rev Mol Cell Biol. 6(6):439-48. (2005) Greenberg JT. Dev Cell. 8(6):799-801. (2005) Levine B. Cell. 120(2):159-62. (2005) Shintani T and Klionsky DJ. Science. 306(5698):990-5. (2004) Tanida I., et al. Int. J. Biochem. Cell Biol. 36:2503-2518(2004) He H., et al. J. Biol. Chem. 278:29278-29287(2003) Tanida I., et al. J. Biol. Chem. 279:36268-36276(2004)

### **Phospho-LC3C(S12) Antibody - Citations**

- [PKC \$\lambda\$ /I Loss Induces Autophagy, Oxidative Phosphorylation, and NRF2 to Promote Liver Cancer Progression](#)
- [Assessment of Posttranslational Modifications of ATG proteins.](#)
- [The endothelial adrenomedullin-RAMP2 system regulates vascular integrity and suppresses tumour metastasis.](#)
- [Identification of a small molecule targeting annexin A7.](#)
- [MAPK15/ERK8 stimulates autophagy by interacting with LC3 and GABARAP proteins.](#)
- [Rab5 and class III phosphoinositide 3-kinase Vps34 are involved in hepatitis C virus NS4B-induced autophagy.](#)
- [Regulation of the autophagy protein LC3 by phosphorylation.](#)