

Phospho-EGFR(Y1069) Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3307a

Specification

Phospho-EGFR(Y1069) Antibody - Product Information

Application	WB, DB,E
Primary Accession	P00533
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	134277

Phospho-EGFR(Y1069) Antibody - Additional Information

Gene ID 1956

Other Names

Epidermal growth factor receptor,
Proto-oncogene c-ErbB-1, Receptor
tyrosine-protein kinase erbB-1, EGFR, ERBB,
ERBB1, HER1

Target/Specificity

This EGFR Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y1069 of human EGFR.

Dilution

WB~~1:2000
DB~~1:500


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



Western blot analysis of lysates from A431 cell line, untreated or treated with EGF, 100ng/ml, using EGFR(Y1069) Antibody (Cat. #AP3307a)(upper) or EGFR (lower).



(LEFT)Dot blot analysis of Phospho-EGFR-Y1069 polyclonal antibody (Cat# AP3307a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentration was 0.5ug per ml. P-Pab: phospho-antibody; P-Peptide: phospho-peptide; NP-Peptide: non-phospho-peptide. (RIGHT)Western blot analysis of EGFR(Y1069) in HeLa cell lysates,

Phospho-EGFR(Y1069) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-EGFR(Y1069) Antibody - Protein Information

Name EGFR ([HGNC:3236](#))

Synonyms ERBB, ERBB1, HER1

Function

Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses (PubMed:<[, PubMed:<\[, PubMed:<\\[\\\). Known ligands include EGF, TGFA/TGF-alpha, AREG, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF \\\(PubMed:<\\\[, PubMed:<\\\\[, PubMed:<\\\\\[, PubMed:<\\\\\\[, PubMed:<\\\\\\\[, PubMed:<\\\\\\\\[, PubMed:<\\\\\\\\\[, PubMed:<\\\\\\\\\\[, PubMed:<\\\\\\\\\\\[\\\\\\\\\\\\). Ligand binding triggers receptor homo- and/or\\\\\\\\\\\]\\\\\\\\\\\(http://www.uniprot.org/citations/17909029\\\\\\\\\\\)\\\\\\\\\\]\\\\\\\\\\(http://www.uniprot.org/citations/20837704\\\\\\\\\\)\\\\\\\\\]\\\\\\\\\(http://www.uniprot.org/citations/27153536\\\\\\\\\)\\\\\\\\]\\\\\\\\(http://www.uniprot.org/citations/12297049\\\\\\\\)\\\\\\\]\\\\\\\(http://www.uniprot.org/citations/15611079\\\\\\\)\\\\\\]\\\\\\(http://www.uniprot.org/citations/9419975\\\\\\)\\\\\]\\\\\(http://www.uniprot.org/citations/8144591\\\\\)\\\\]\\\\(http://www.uniprot.org/citations/7679104\\\\)\\\]\\\(http://www.uniprot.org/citations/2790960\\\)\\]\\(http://www.uniprot.org/citations/27153536\\)\]\(http://www.uniprot.org/citations/10805725\)](http://www.uniprot.org/citations/2790960)

either noninduced (Lane 1) or induced with EGF (Lane 2).

Phospho-EGFR(Y1069) Antibody - Background

Epidermal Growth factor receptor (EGFR) is the prototype member of the type 1 receptor tyrosine kinases. EGFR overexpression in tumors indicates poor prognosis and is observed in tumors of the head and neck, brain, bladder, stomach, breast, lung, endometrium, cervix, vulva, ovary, esophagus, stomach and in squamous cell carcinoma. EGFR is a receptor for EGF, but also for other members of the EGF family, including TGF-alpha, amphiregulin, betacellulin, heparin-binding EGF-like growth factor, GP30 and vaccinia virus growth factor. Is involved in the control of cell growth and differentiation.

Phospho-EGFR(Y1069) Antibody - References

Aifa, S., et al., Exp. Cell Res. 302(1):108-114 (2005).
Adams, T.E., et al., Growth Factors 22(2):89-95 (2004).
Ichinose, J., et al., Biochem. Biophys. Res. Commun. 324(3):1143-1149 (2004).
Kuribayashi, A., et al., Endocrinology 145(11):4976-4984 (2004).
Kapoor, G.S., et al., Mol. Cell. Biol. 24(2):823-836 (2004).

heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules (PubMed:27153536). May also activate the NF-kappa-B signaling cascade (PubMed:11116146). Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling (PubMed:11602604). Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin (PubMed:11483589). Positively regulates cell migration via interaction with CCDC88A/GIV which retains EGFR at the cell membrane following ligand stimulation, promoting EGFR signaling which triggers cell migration (PubMed:20462955). Plays a role in enhancing learning and memory performance (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein. Golgi apparatus membrane; Single-pass type I membrane protein. Nucleus membrane; Single-pass type I membrane protein Endosome Endosome membrane. Nucleus. Note=In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER (PubMed:20674546, PubMed:17909029). Endocytosed upon activation by ligand (PubMed:2790960, PubMed:17182860, PubMed:27153536, PubMed:17909029). Colocalized with GPER1 in the nucleus of estrogen agonist-induced cancer-associated fibroblasts (CAF) (PubMed:20551055)

Tissue Location

Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.

**Phospho-EGFR(Y1069) Antibody -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)