

Phospho-MAP3K7(T187) Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3323a

Specification

Phospho-MAP3K7(T187) Antibody - Product Information

Application	DB,E
Primary Accession	Q43318
Other Accession	P0C8E4 , Q62073 , A2VDU3
Reactivity Predicted	Human Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	67196

Phospho-MAP3K7(T187) Antibody - Additional Information

Gene ID 6885

Other Names

Mitogen-activated protein kinase kinase 7, Transforming growth factor-beta-activated kinase 1, TGF-beta-activated kinase 1, MAP3K7, TAK1

Target/Specificity

This MAP3K7 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding T187 of human MAP3K7.

Dilution


DB~~1:500

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw



Dot blot analysis of Phospho-MAP3K7-T187 polyclonal antibody (Cat.AP3323a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentration were 0.5ug per ml.

Phospho-MAP3K7(T187) Antibody - Background

MAP3K7 is a member of the serine/threonine protein kinase family. This kinase mediates the signaling transduction induced by TGF beta and morphogenetic protein (BMP), and controls a variety of cell functions including transcription regulation and apoptosis. In response to IL-1, this protein forms a kinase complex including TRAF6, MAP3K7P1/TAB1 and MAP3K7P2/TAB2; this complex is required for the activation of nuclear factor kappa B. This kinase can also activate MAPK8/JNK, MAP2K4/MKK4, and thus plays a role in the cell response to environmental stresses.

Phospho-MAP3K7(T187) Antibody - References

Smit, L., et al., J. Biol. Chem. 279(17):17232-17240 (2004).

cycles.

Precautions

Phospho-MAP3K7(T187) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-MAP3K7(T187) Antibody - Protein Information

Name MAP3K7

{ECO:0000303|PubMed:28397838,
ECO:0000312|HGNC:HGNC:6859}

Function

Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. Plays an important role in the cascades of cellular responses evoked by changes in the environment. Mediates signal transduction of TRAF6, various cytokines including interleukin-1 (IL-1), transforming growth factor-beta (TGFB), TGFB-related factors like BMP2 and BMP4, toll-like receptors (TLR), tumor necrosis factor receptor CD40 and B-cell receptor (BCR). Ceramides are also able to activate MAP3K7/TAK1. Once activated, acts as an upstream activator of the MKK/JNK signal transduction cascade and the p38 MAPK signal transduction cascade through the phosphorylation and activation of several MAP kinase kinases like MAP2K1/MEK1, MAP2K3/MKK3, MAP2K6/MKK6 and MAP2K7/MKK7. These MAP2Ks in turn activate p38 MAPKs, c-jun N-terminal kinases (JNKs) and I-kappa-B kinase complex (IKK). Both p38 MAPK and JNK pathways control the transcription factors activator protein-1 (AP-1), while nuclear factor-kappa B is activated by IKK. MAP3K7 activates also IKBKB and MAPK8/JNK1 in response to TRAF6 signaling and mediates BMP2-induced apoptosis. In osmotic stress signaling, plays a major role in the activation of MAPK8/JNK1, but not that of NF-kappa-B. Promotes TRIM5 capsid-specific restriction activity. Phosphorylates RIPK1 at 'Ser-321' which positively regulates RIPK1 interaction with RIPK3 to promote necroptosis but negatively regulates RIPK1 kinase activity and its interaction with FADD to mediate apoptosis (By similarity).

Cellular Location

Ono, K., et al., Biochem. Biophys. Res. Commun. 307(2):332-337 (2003).
Sakurai, H., et al., J. Biol. Chem. 278(38):36916-36923 (2003).
Cheung, P.C., et al., EMBO J. 22(21):5793-5805 (2003).
Edlund, S., et al., Mol. Biol. Cell 14(2):529-544 (2003).

Cytoplasm. Cell membrane; Peripheral membrane protein; Cytoplasmic side.
Note=Although the majority of MAP3K7/TAK1 is found in the cytosol, when complexed with TAB1/MAP3K7IP1 and TAB2/MAP3K7IP2, it is also localized at the cell membrane

Tissue Location

Isoform 1A is the most abundant in ovary, skeletal muscle, spleen and blood mononuclear cells. Isoform 1B is highly expressed in brain, kidney and small intestine. Isoform 1C is the major form in prostate. Isoform 1D is the less abundant form

Phospho-MAP3K7(T187) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-MAP3K7(T187) Antibody - Citations

- [Targeting of TGF- \$\beta\$ -activated protein kinase 1 inhibits chemokine \(C-C motif\) receptor 7 expression, tumor growth and metastasis in breast cancer.](#)