

**Phospho-MAP4K4(S629) Antibody**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP3428a**

**Specification**

**Phospho-MAP4K4(S629) Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | DB,E                   |
| Primary Accession | <a href="#">O95819</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Isotype           | Rabbit IgG             |

**Phospho-MAP4K4(S629) Antibody - Additional Information**

**Gene ID** 9448

**Other Names**

Mitogen-activated protein kinase kinase kinase 4, HPK/GCK-like kinase HGK, MAPK/ERK kinase kinase kinase 4, MEK kinase kinase 4, MEKKK 4, Nck-interacting kinase, MAP4K4, HGK, KIAA0687, NIK

**Target/Specificity**

This Phospho-MAP4K4-pS629 antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S629 of human MAP4K4.

**Dilution**

DB~~1:500

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Phospho-MAP4K4(S629) Antibody is for



Dot blot analysis of anti-MAP4K4-pS629 Phospho-specific Pab (Cat.#AP3428a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

**Phospho-MAP4K4(S629) Antibody - Background**

M4K4, a member of the STE20 subfamily of Ser/Thr protein kinases, may play a role in the response to environmental stress and cytokines such as TNF-alpha. It appears to act upstream of the JUN N-terminal pathway. This protein is thought to interact with the SH3 domain of the adapter proteins Nck. HGK binds, via its CNH regulatory domain, to the N-terminal region of SPG3A. Expression appears to be ubiquitous, expressed in all tissue types examined. Isoform 5 appears to be more abundant in the brain, and isoform 4 is predominant in the liver, skeletal muscle and placenta.

**Phospho-MAP4K4(S629) Antibody - References**

Wright, J.H., et al., Mol. Cell. Biol.

research use only and not for use in diagnostic or therapeutic procedures.

**Phospho-MAP4K4(S629) Antibody - Protein Information**

**Name** MAP4K4

**Synonyms** HGK, KIAA0687, NIK

**Function**

Serine/threonine kinase that may play a role in the response to environmental stress and cytokines such as TNF-alpha. Appears to act upstream of the JUN N-terminal pathway. Phosphorylates SMAD1 on Thr-322.

**Cellular Location**

Cytoplasm.

**Tissue Location**

Widely expressed. Isoform 5 is abundant in the brain. Isoform 4 is predominant in the liver, skeletal muscle and placenta.

23(6):2068-2082 (2003).

Yao, Z., et al., J. Biol. Chem. 274(4):2118-2125 (1999).

Ishikawa, K., et al., DNA Res. 5(3):169-176 (1998).

**Phospho-MAP4K4(S629) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)