

Phospho-LC3B(T12) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP3530a

Specification

Phospho-LC3B(T12) Antibody - Product Information

Application	DB,E
Primary Accession	Q9GZQ8
Other Accession	A6NCE7 , O41515
Reactivity	Human
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig

Phospho-LC3B(T12) Antibody - Additional Information

Gene ID 81631

Other Names

Microtubule-associated proteins 1A/1B light chain 3B, Autophagy-related protein LC3 B, Autophagy-related ubiquitin-like modifier LC3 B, MAP1 light chain 3-like protein 2, MAP1A/MAP1B light chain 3 B, MAP1A/MAP1B LC3 B, Microtubule-associated protein 1 light chain 3 beta, MAP1LC3B, MAP1ALC3

Target/Specificity

This LC3B Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding T12 of human LC3B.

Dilution


DB~~1:500

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C



Dot blot analysis of Phospho-LC3 (APG8b)-T12 Antibody (Cat.#AP3530a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

Phospho-LC3B(T12) Antibody - Background

MAP1A and MAP1B are microtubule-associated proteins which mediate the physical interactions between microtubules and components of the cytoskeleton. These proteins are involved in formation of autophagosomal vacuoles (autophagosomes). MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. MAP1LC3b is one of the light chain subunits and can associate with either MAP1A or MAP1B. The precursor molecule is cleaved by APG4B/ATG4B to form the cytosolic form, LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form the membrane-bound form, LC3-II.

Macroautophagy is the major inducible

in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-LC3B(T12) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-LC3B(T12) Antibody - Protein Information

Name MAP1LC3B ([HGNC:13352](#))

Synonyms MAP1ALC3

Function

Ubiquitin-like modifier involved in formation of autophagosomal vacuoles (autophagosomes). Plays a role in mitophagy which contributes to regulate mitochondrial quantity and quality by eliminating the mitochondria to a basal level to fulfill cellular energy requirements and preventing excess ROS production (PubMed:<[a href="http://www.uniprot.org/citations/28017329"](http://www.uniprot.org/citations/28017329)target="_blank">28017329). While LC3s are involved in elongation of the phagophore membrane, the GABARAP/GATE-16 subfamily is essential for a later stage in autophagosome maturation. Promotes primary ciliogenesis by removing OFD1 from centriolar satellites via the autophagic pathway. Through its interaction with the reticulophagy receptor TEX264, participates in the remodeling of subdomains of the endoplasmic reticulum into autophagosomes upon nutrient stress, which then fuse with lysosomes for endoplasmic reticulum turnover (PubMed:<[a href="http://www.uniprot.org/citations/31006538"](http://www.uniprot.org/citations/31006538)target="_blank">31006538, PubMed:<[a href="http://www.uniprot.org/citations/31006537"](http://www.uniprot.org/citations/31006537)target="_blank">31006537).

Cellular Location

Cytoplasm, cytoskeleton. Endomembrane system; Lipid-anchor. Cytoplasmic vesicle, autophagosome membrane; Lipid-anchor. Note=LC3-II binds to the autophagic membranes. LC3-II localizes with the mitochondrial inner membrane during Parkin-mediated mitophagy (PubMed:28017329). Localizes also to

pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole).

Phospho-LC3B(T12) Antibody - References

Baehrecke EH. Nat Rev Mol Cell Biol. 6(6):505-10. (2005)
Lum JJ, et al. Nat Rev Mol Cell Biol. 6(6):439-48. (2005)
Greenberg JT. Dev Cell. 8(6):799-801. (2005)
Levine B. Cell. 120(2):159-62. (2005)
Shintani T and Klionsky DJ. Science. 306(5698):990-5. (2004)
Tanida I., et al. Int. J. Biochem. Cell Biol. 36:2503-2518(2004)
He H., et al. J. Biol. Chem. 278:29278-29287(2003)
Tanida I., et al. J. Biol. Chem. 279:36268-36276(2004)

discrete punctae along the ciliary axoneme

Tissue Location

Most abundant in heart, brain, skeletal muscle and testis. Little expression observed in liver

Phospho-LC3B(T12) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-LC3B(T12) Antibody - Citations

- [Regulation of the autophagic machinery in human neutrophils.](#)