

### Bi-Phospho-APG8b(MAP1LC3B)(T93/Y99) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP3739a

### **Specification**

#### Bi-Phospho-APG8b(MAP1LC3B)(T93/Y99) Antibody - Product Information

Application
Primary Accession
Other Accession

Other Accession

DB,E

Q9GZQ8

A6NCE7,

NP\_073729

Reactivity Human
Host Rabbit
Clonality Polyclonal
Isotype Rabbit Ig
Calculated MW 14688

Bi-Phospho-APG8b(MAP1LC3B)(T93/Y99) Antibody - Additional Information

#### Gene ID 81631

#### **Other Names**

Microtubule-associated proteins 1A/1B light chain 3B, Autophagy-related protein LC3 B, Autophagy-related ubiquitin-like modifier LC3 B, MAP1 light chain 3-like protein 2, MAP1A/MAP1B light chain 3 B, MAP1A/MAP1B LC3 B, Microtubule-associated protein 1 light chain 3 beta, MAP1LC3B, MAP1ALC3

## **Target/Specificity**

This APG8b(MAP1LC3B) Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding T93/Y99 of human APG8b(MAP1LC3B).


### Dilution

DB~~1:500

### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

### Storage




Dot blot analysis of anti-Phospho-APG8b (MAP1LC3B)-T93/Y99 Phospho-specific Pab (Cat. #AP3739a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

## Bi-Phospho-APG8b(MAP1LC3B)(T93/Y99) Antibody - Background

The product of this gene is a subunit of neuronal microtubule-associated MAP1A and MAP1B proteins, which are involved in microtubule assembly and important for neurogenesis. Studies on the rat homolog implicate a role for this gene in autophagy, a process that involves the bulk degradation of cytoplasmic component.

# Bi-Phospho-APG8b(MAP1LC3B)(T93/Y99) Antibody - References

Rouschop, K.M., et al. J. Clin. Invest. 120(1):127-141(2010) Kirkin, V., et al. Mol. Cell 33(4):505-516(2009)





Tel: 858.875.1900 Fax: 858.622.0609

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

Bi-Phospho-APG8b(MAP1LC3B)(T93/Y99) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Bi-Phospho-APG8b(MAP1LC3B)(T93/Y99) **Antibody - Protein Information** 

Name MAP1LC3B (HGNC:13352)

**Synonyms MAP1ALC3** 


## **Function**

Ubiquitin-like modifier involved in formation of autophagosomal vacuoles (autophagosomes). Plays a role in mitophagy which contributes to regulate mitochondrial quantity and quality by eliminating the mitochondria to a basal level to fulfill cellular energy requirements and preventing excess ROS production (PubMed:<a href="http://www.uniprot.org/c itations/28017329" target="\_blank">28017329</a>). While LC3s are involved in elongation of the phagophore membrane, the GABARAP/GATE-16 subfamily is essential for a later stage in autophagosome maturation. Promotes primary ciliogenesis by removing OFD1 from centriolar satellites via the autophagic pathway. Through its interaction with the reticulophagy receptor TEX264, participates in the remodeling of subdomains of the endoplasmic reticulum into autophagosomes upon nutrient stress, which then fuse with lysosomes for endoplasmic reticulum turnover (PubMed:<a href="http://www.uniprot.org/c itations/31006538" target="\_blank">31006538</a>, PubMed:<a href="http://www.uniprot.org/ci tations/31006537" target=" blank">31006537</a>).

## **Cellular Location**

Cytoplasm, cytoskeleton. Endomembrane system; Lipid-anchor. Cytoplasmic vesicle, autophagosome membrane; Lipid-anchor. Note=LC3-II binds to the autophagic membranes. LC3-II localizes with the

Othman, E.Q., et al. J. Clin. Lab. Anal. 23(4):249-258(2009) Liu, Q., et al. Ai Zheng 27(1):25-29(2008) Komatsu, M., et al. Cell 131(6):1149-1163(2007)





mitochondrial inner membrane during Parkin-mediated mitophagy (PubMed:28017329). Localizes also to discrete punctae along the ciliary axoneme

## **Tissue Location**

Most abundant in heart, brain, skeletal muscle and testis. Little expression observed in liver

# Bi-Phospho-APG8b(MAP1LC3B)(T93/Y99) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture