

Phospho-ULK1(S317) Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3803a

Specification

Phospho-ULK1(S317) Antibody - Product Information

Application	DB,E
Primary Accession	O75385
Other Accession	NP_003556.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig

Phospho-ULK1(S317) Antibody - Additional Information

Gene ID 8408

Other Names

Serine/threonine-protein kinase ULK1,
Autophagy-related protein 1 homolog,
ATG1, hATG1, Unc-51-like kinase 1, ULK1,
KIAA0722

Target/Specificity

This ULK1 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S317 of human ULK1.

Dilution

DB~~1:500

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-ULK1(S317) Antibody is for



Dot blot analysis of ULK1 Antibody (Phospho S317) Phospho-specific Pab (Cat. #AP3803a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.6ug per ml.

Phospho-ULK1(S317) Antibody - Background

Involved in autophagy. Required for autophagosome formation (By similarity). Target of the TOR kinase signaling pathway that regulates autophagy through the control of phosphorylation status of ATG13/KIAA0652 and ULK1, and the regulation of the ATG13-ULK1-RB1CC1 complex (By similarity). Phosphorylates ATG13/KIAA0652. Involved in axon growth (By similarity). Plays an essential role in neurite extension of cerebellar granule cells (By similarity).

Phospho-ULK1(S317) Antibody - References

Mercer, C.A., et al. Autophagy 5(5):649-662(2009) Ganley, I.G., et al. J. Biol. Chem. 284(18):12297-12305(2009) Jung, C.H., et al. Mol. Biol. Cell 20(7):1992-2003(2009) Hosokawa, N., et al. Mol. Biol. Cell 20(7):1981-1991(2009) Chan, E.Y. Sci Signal 2 (84), PE51 (2009) :

research use only and not for use in
diagnostic or therapeutic procedures.

Phospho-ULK1(S317) Antibody - Protein Information

Name ULK1 ([HGNC:12558](#))

Synonyms KIAA0722

Function

Serine/threonine-protein kinase involved in autophagy in response to starvation (PubMed:[18936157](http://www.uniprot.org/citations/18936157)), PubMed:[21460634](http://www.uniprot.org/citations/21460634), PubMed:[21795849](http://www.uniprot.org/citations/21795849), PubMed:[25040165](http://www.uniprot.org/citations/25040165)). Acts upstream of phosphatidylinositol 3-kinase PIK3C3 to regulate the formation of autophagophores, the precursors of autophagosomes (PubMed:[18936157](http://www.uniprot.org/citations/18936157)), PubMed:[21460634](http://www.uniprot.org/citations/21460634), PubMed:[21795849](http://www.uniprot.org/citations/21795849), PubMed:[25040165](http://www.uniprot.org/citations/25040165)). Part of regulatory feedback loops in autophagy: acts both as a downstream effector and negative regulator of mammalian target of rapamycin complex 1 (mTORC1) via interaction with RPTOR (PubMed:[21795849](http://www.uniprot.org/citations/21795849)). Activated via phosphorylation by AMPK and also acts as a regulator of AMPK by mediating phosphorylation of AMPK subunits PRKAA1, PRKAB2 and PRKAG1, leading to negatively regulate AMPK activity (PubMed:[21460634](http://www.uniprot.org/citations/21460634)). May phosphorylate ATG13/KIAA0652 and RPTOR;

however such data need additional evidences (PubMed:18936157). Plays a role early in neuronal differentiation and is required for granule cell axon formation (PubMed:11146101). May also phosphorylate SESN2 and SQSTM1 to regulate autophagy (PubMed:25040165). Phosphorylates FLCN, promoting autophagy (PubMed:25126726).

Cellular Location

Cytoplasm, cytosol. Preautophagosomal structure. Note=Under starvation conditions, is localized to punctate structures primarily representing the isolation membrane that sequesters a portion of the cytoplasm resulting in the formation of an autophagosome.

Tissue Location

Ubiquitously expressed. Detected in the following adult tissues: skeletal muscle, heart, pancreas, brain, placenta, liver, kidney, and lung

Phospho-ULK1(S317) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-ULK1(S317) Antibody - Citations

- [GZ17-6.02 and Doxorubicin Interact to Kill Sarcoma Cells via Autophagy and Death Receptor Signaling](#)
- [Neratinib decreases pro-survival responses of \[sorafenib + vorinostat\] in pancreatic cancer](#)
- [The multi-kinase inhibitor lenvatinib interacts with the HDAC inhibitor entinostat to kill liver cancer cells](#)
- [Enhanced signaling via ERBB3/PI3K plays a compensatory survival role in pancreatic tumor cells exposed to \[neratinib + valproate\]](#)

- [Fingolimod Augments Monomethylfumarate Killing of GBM Cells](#)
- [\(Curcumin+sildenafil\) enhances the efficacy of 5FU and anti-PD1 therapies in vivo](#)
- [GZ17-6.02 initiates DNA damage causing autophagosome-dependent HDAC degradation resulting in enhanced anti-PD1 checkpoint inhibitory antibody efficacy](#)