

Phospho-mouse p21Cip1(S78) Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3877a

Specification

Phospho-mouse p21Cip1(S78) Antibody - Product Information

Application	DB,E
Primary Accession	P39689
Other Accession	NP_001129489.1
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	17785

Phospho-mouse p21Cip1(S78) Antibody - Additional Information

Gene ID 12575

Other Names

Cyclin-dependent kinase inhibitor 1, CDK-interacting protein 1, Melanoma differentiation-associated protein, p21, Cdkn1a, Cip1, Waf1

Target/Specificity

This mouse p21Cip1 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S78 of mouse p21Cip1.

Dilution

DB~~1:500


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



Dot blot analysis of Mouse p21Cip1 Antibody (Phospho S78) Phospho-specific Pab (Cat. #AP3877a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.6ug per ml.

Phospho-mouse p21Cip1(S78) Antibody - Background

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be

Phospho-mouse p21Cip1(S78) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-mouse p21Cip1(S78) Antibody - Protein Information

Name Cdkn1a

Synonyms Cip1, Waf1

Function

May be involved in p53/TP53 mediated inhibition of cellular proliferation in response to DNA damage. Binds to and inhibits cyclin- dependent kinase activity, preventing phosphorylation of critical cyclin-dependent kinase substrates and blocking cell cycle progression. Functions in the nuclear localization and assembly of cyclin D-CDK4 complex and promotes its kinase activity towards RB1. At higher stoichiometric ratios, inhibits the kinase activity of the cyclin D- CDK4 complex (PubMed:25329316). Inhibits DNA synthesis by DNA polymerase delta by competing with POLD3 for PCNA binding (By similarity). Plays an important role in controlling cell cycle progression and DNA damage-induced G2 arrest (By similarity).

Cellular Location

Cytoplasm
{ECO:0000250|UniProtKB:P38936}. Nucleus

involved in the phosphorylation of tumor suppressor protein Rb. The CDK4 activity associated with this cyclin was reported to be necessary for cell cycle progression through G2 phase into mitosis after UV radiation. Several transcript variants encoding different isoforms have been found for this gene.

Phospho-mouse p21Cip1(S78) Antibody - References

Liu, C.Y., et al. Carcinogenesis 31(7):1259-1263(2010)
Kim, J., et al. Cytokine 50(1):42-49(2010)
Kamatani, Y., et al. Nat. Genet. 42(3):210-215(2010)
Gumina, M.R., et al. Cell Cycle 9(4):820-828(2010)
Radulovich, N., et al. Mol. Cancer 9, 24 (2010) :

Phospho-mouse p21Cip1(S78) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)